

Please simplify your answers to the extent reasonable without a calculator. Show your work. Explain your answers, concisely. In case you need them, here are the probability distributions we have learned ($0 \leq p \leq 1$, $0 < n \in \mathbb{Z}$, $0 < \lambda, \sigma \in \mathbb{R}$, $a < b \in \mathbb{R}$, $\mu \in \mathbb{R}$, $1 \leq r \in \mathbb{R}$, and any values not listed for the random variables have probability 0 or probability density 0):

$$\begin{aligned}
 B \sim \text{Ber}(p) & \quad P(B = b) = \begin{cases} 1 - p & \text{if } b = 0; \\ p & \text{if } b = 1. \end{cases} \\
 K \sim \text{Bin}(n, p) & \quad P(K = k) = \binom{n}{k} p^k (1 - p)^{n-k}, \quad k \in \{0, 1, \dots, n\}. \\
 K \sim \text{Geom}(p) & \quad P(K = k) = (1 - p)^{k-1} p, \quad 0 < k \in \mathbb{Z}. \\
 K \sim \text{Poisson}(\lambda) & \quad P(K = k) = \frac{\lambda^k}{k!} e^{-\lambda}, \quad 0 \leq k \in \mathbb{Z}. \\
 X \sim \text{Unif}[a, b] & \quad f(x) = \frac{1}{b - a}, \quad x \in [a, b]. \\
 X \sim \mathcal{N}(\mu, \sigma^2) & \quad f(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-(x-\mu)^2/(2\sigma^2)}, \quad x \in \mathbb{R}. \\
 T \sim \text{Exp}(\lambda) & \quad f(t) = \lambda e^{-\lambda t}, \quad 0 \leq t \in \mathbb{R}. \\
 T \sim \text{Gamma}(r, \lambda) & \quad f(t) = \frac{\lambda^r t^{r-1}}{\Gamma(r)} e^{-\lambda t}, \quad 0 \leq t \in \mathbb{R}.
 \end{aligned}$$

1. [25 points] For $1 < \alpha \in \mathbb{R}$, let X_α be a continuous random variable with probability density function:

$$f_\alpha(x) = \begin{cases} (\alpha - 1)x^{-\alpha} & \text{if } 1 \leq x \in \mathbb{R}; \\ 0 & \text{otherwise.} \end{cases}$$

- a. [5 points] Show that f_α is a probability density function for $\alpha > 1$.
 - b. [10 points] For which values of α is $\mathbf{E}[X_\alpha] \in \mathbb{R}$? For these values of α , what is $\mathbf{E}[X_\alpha]$?
 - c. [10 points] For which values of α is $\mathbf{Var}[X_\alpha] \in \mathbb{R}$? For these values of α , what is $\mathbf{Var}[X_\alpha]$?
2. Human heights have approximately normal distributions: American women with a mean of about 64 inches and a standard deviation of 2.5 inches; American men with a mean of about 69.5 inches and a standard deviation of 3 inches.*
- a. [8 points] Explain why a normal distribution can't be exactly right for human heights, but could still be a good approximation.
 - b. [7 points] Mary is one standard deviation taller than the average American woman. Approximately what fraction of American women is she taller than?
 - c. [10 points] Approximately what fraction of American men is Mary taller than?

* See M. F. Schilling, A. E. Watkins and W. Watkins, "Is human height bimodal?", *The American Statistician* **56** (2012) 223–229.

3. [25 points] Recall that a Poisson process with intensity λ is defined to be a set of random points on $[0, \infty)$, satisfying three properties: the points are distinct; for a bounded interval $I \subset [0, \infty)$ the number of points $N(I) \sim \text{Poisson}(\lambda|I|)$; and for non-overlapping intervals I_1, \dots, I_n , the random variables $N(I_1), \dots, N(I_n)$ are mutually independent.
 - a. [9 points] Argue that for any $a > 0$, the points of a Poisson process with intensity λ on $[0, \infty)$, that lie in the interval $[a, \infty)$, satisfy the same three properties.
 - b. [16 points] Let T_1 be the location of the first (smallest coordinate) point in a Poisson process with intensity $\lambda > 0$ on $[0, \infty)$. Let T_2 be the location of the second point. Use the result of part (a) to find the probability density function for $T_2 - T_1$.
4. [25 points] Let $Z \sim \mathcal{N}(0, 1)$ be a standard normal random variable. Find the moment generating function $M(t)$ of Z .