

Math 203, Problem Set 4. Due Wednesday, November 25.

For this problem set, you may assume that the ground field is algebraically closed.

1. (*Hyperelliptic curves.*) Let a_1, \dots, a_5 be pairwise distinct constants. Find the singularities of the projective hyperelliptic curve of genus 2:

$$y^2 z^3 = (x - a_1 z) \dots (x - a_5 z).$$

2. (*Pencils of conics and singularities.*) Let Q_1 and Q_2 be two distinct nonsingular conics in \mathbb{P}^2 . The family of conics

$$Q_{\lambda, \mu} = \lambda Q_1 + \mu Q_2$$

where $[\lambda : \mu] \in \mathbb{P}^1$ is called a *pencil* of conics.

- (i) Recall that any conic $Q \subset \mathbb{P}^2$ determines and is determined by the symmetric matrix A of coefficients with

$$Q([x : y : z]) = [x \ y \ z] A [x \ y \ z]^T.$$

Possibly by diagonalizing A (and therefore Q), show that

$$Q \text{ is singular if and only if } \det A = 0.$$

- (ii) Letting $A_{\lambda, \mu}$ be the matrix associated to the conic $Q_{\lambda, \mu}$, show that $\det A_{\lambda, \mu}$ is a cubic polynomial in λ, μ . Prove that any pencil of conics contains (at most) 3 singular conics.
- (iii) Let p_1, p_2, p_3, p_4 be points in \mathbb{P}^2 such that no three of them lie on a line. Show that the set of conics through p_1, p_2, p_3, p_4 is a pencil. (Feel free to change coordinates to prove this fact). What are the singular conics in this pencil? Can you draw them?

3. (*Singularities of cubics.*)

- (i) Show that any singular irreducible cubic in \mathbb{P}^2 is isomorphic to either the nodal or the cuspidal cubics:

$$y^2 z = x^2(x + z) \text{ or } y^2 z = x^3.$$

Hint: Assume the singularity is at $[0 : 0 : 1]$. Show that the cubic can be written as

$$(\text{quadratic polynomial in } x, y) \cdot z = Q(x, y),$$

where Q is a cubic polynomial in x, y . Change coordinates suitably and write the cubic as

$$y^2 z = \tilde{Q}(x, y) \text{ or } xyz = \tilde{Q}(x, y).$$

Use the coordinate change $z \mapsto \lambda x + \mu y + \nu z$ to put the cubic into one of the forms

$$y^2 z = (x + by)^3 \text{ or } xyz = (x + y)^3.$$

Conclude by performing one more change of coordinates.

- (ii) Using (i), show that *irreducible cubics* in \mathbb{P}^2 can have at most 1 singular point. Exhibit a cubic in \mathbb{P}^2 with 3 singular points.

Remark: It can be shown that an *irreducible* degree d curve in \mathbb{P}^2 has at most $\binom{d-1}{2}$ singular points.

4. Show that a *general* hypersurface of degree d in \mathbb{P}^n is non-singular:

- (i) For any hypersurface $Z(f) \subset \mathbb{P}^n$ of degree d , view the coefficients of f as a point p_f in a large dimensional projective space \mathbb{P}^N (This projective space is called *the moduli space* of degree d hypersurfaces). Let

$$X = \{(f, p) \in \mathbb{P}^N \times \mathbb{P}^n : p \text{ is a singular point of } f\}.$$

Show that X is a projective algebraic set in $\mathbb{P}^N \times \mathbb{P}^n$.

- (ii) Conclude that the image $\pi(X)$ of X under the projection onto \mathbb{P}^N is a projective algebraic set. What is $\pi(X)$? Conclude that the subset of \mathbb{P}^N corresponding to smooth hypersurfaces is open and *nonempty*.

Remark: This will prove in particular that f is singular provided that the coefficients of f satisfy certain polynomial relations. Therefore, if you pick your coefficients of f randomly, these polynomial relations will most likely not be satisfied and your hypersurface is non-singular. This is the explanation of the word *general*.

5. (*Analytic singularities.*) Consider the singular plane curves Z and W given by the equations

$$y^2 - x^2(x + 1) = 0 \text{ and } xy = 0$$

respectively.

- (i) Explain briefly why Z and W are not isomorphic. Explain that $(0, 0)$ is an ordinary double point for both of these curves. What are the tangent directions at $(0, 0)$ for Z and W ? Sketch (the real points of) Z and W . Do Z and W look *alike* near the origin?
- (ii) Show that there are *formal power series*

$$\tilde{x} = f_1 + f_2 + f_3 + \dots \text{ and}$$

$$\tilde{y} = g_1 + g_2 + g_3 \dots$$

in the variables x and y such that the equation of Z becomes

$$\tilde{x}\tilde{y} = 0.$$

Hint: Construct the degree i homogeneous parts f_i and g_i inductively. Show you can pick

$$f_1 = y - x, g_1 = x + y.$$

Next, you would need

$$f_2(x + y) + g_2(y - x) = -x^3.$$

Why can you construct f_2 and g_2 ? Continue in this fashion.

Remark: If we work over an arbitrary field k it doesn't make sense to ask if the power series \tilde{x} and \tilde{y} converge, hence the terminology *formal power series*. Convergence may be arranged if you work over the complex numbers, but you don't have to prove it.

Remark: It turns out the assignment

$$(x, y) \rightarrow (\tilde{x}, \tilde{y})$$

is invertible *e.g.* you can solve for x, y in terms of formal power series in \tilde{x}, \tilde{y} . In fact, this statement is generally true about any power series

$$\tilde{x} = ax + by + \dots, \tilde{y} = cx + dy + \dots$$

provided that $ad - bc \neq 0$. Therefore, the assignment

$$(x, y) \rightarrow (\tilde{x}, \tilde{y})$$

is a *formal* change of coordinates, establishing a *formal isomorphism* between Z and W . We say that Z and W are *analytically equivalent*.

Remark: Over the complex numbers, convergence may be arranged near the origin, if x, y are small, and thus the word *formal* may be replaced by *local analytic isomorphism* near the origin.

- (iii) Explain briefly why any ordinary double point singularity in \mathbb{A}^2 is analytically equivalent to the node $\tilde{x}\tilde{y} = 0$.

Remark: It can be shown that any double point is analytically equivalent to the singularity $\tilde{y}^2 = \tilde{x}^r$, for some r . The case $r = 2$ corresponds to the case which concerned us above.

6. (*Quadrics are rational.*) We have seen in class that an irreducible projective quadric Q corresponds to a symmetric square matrix A such that

$$Q(\mathbf{x}) = \mathbf{x}^T A \mathbf{x} \text{ for } \mathbf{x} \in \mathbb{P}^n.$$

We say that the quadric Q is nondegenerate if the matrix A is non-degenerate.

- (i) Show that a non-degenerate irreducible quadric Q in \mathbb{P}^3 can be written in the form

$$xy = zw$$

after a suitable change of homogeneous coordinates. Conclude that Q is isomorphic to $\mathbb{P}^1 \times \mathbb{P}^1$, hence Q is rational.

- (ii) In general, show that any non-degenerate irreducible quadric $Q \subset \mathbb{P}^{n+1}$ is rational.

Hint: Pick $p \in Q$ and assume after a change of coordinates that $p = [1 : 0 : \dots : 0]$. Consider the linear hyperplane

$$H = \{X_0 = 0\}.$$

For any $q \in Q$, define $f(q)$ to be the point of intersection of the line pq with H . Show that

$$f : Q \dashrightarrow H$$

is a birational isomorphism. This rational map is called the linear projection from p .