

Math 203, Problem Set 6. Due Friday, March 9.

1. (*Affine morphisms.*) Let $f : X \rightarrow Y$ be an affine morphism between two Noetherian separated schemes, and let $\mathcal{F} \rightarrow X$ be a quasicoherent sheaf. Show that

$$H^i(X, \mathcal{F}) = H^i(Y, f_*\mathcal{F}).$$

Hint: Consider the affine open cover \mathcal{U} of Y , and the associated cover $f^{-1}(\mathcal{U})$ for X .

2. (*Hilbert polynomials and arithmetic genus.*) Let $\mathcal{F} \rightarrow X$ be a coherent sheaf over a projective variety $X \subset \mathbb{P}_k^r$.

(i) Show that there exists a polynomial $\chi_{\mathcal{F}}$ such that

$$\chi(\mathcal{F}(n)) = \chi_{\mathcal{F}}(n).$$

Hint: First reduce to the case $X = \mathbb{P}^r$ by considering $\iota_*\mathcal{F}$. Over \mathbb{P}^r , you may wish to argue by induction on r , using the exact sequence

$$0 \rightarrow \mathcal{F}(-H) \rightarrow \mathcal{F} \rightarrow \mathcal{F}|_H \rightarrow 0$$

for a suitable hyperplane $H \subset \mathbb{P}^r$. The only issue is exactness on the left. You can construct a suitable hyperplane working over affine patches and using a bit of commutative algebra (associated points).

(ii) In particular, if $\mathcal{F} = \mathcal{O}_X$ this recovers the Hilbert polynomial we introduced in Math 203a.

(iii) Calculate the Hilbert polynomial of $\mathcal{O}_{\mathbb{P}^r}(m)$.

(iv) Using (ii), show that the arithmetic genus of X is defined as

$$p_a(X) = (-1)^{\dim X - 1}(\chi(\mathcal{O}_X) - 1).$$

In particular, if X is an irreducible projective curve then

$$p_a(X) = \dim H^1(X, \mathcal{O}_X).$$

This definition is independent of the projective embedding.

3. (*Complete intersections.*) Let $X \subset \mathbb{P}^r$ be a smooth complete intersection in projective space with $\dim X = d$. Show that

(i) for all ℓ , the map

$$H^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(\ell)) \rightarrow H^0(X, \mathcal{O}_X(\ell))$$

is surjective. This was stated as a fact in a previous problem set, but now you have the tools to construct a proof.

(ii) In particular, setting $\ell = 0$, X is connected.

(iii) Show that the intermediate cohomology of the twisting line bundles vanishes

$$H^i(X, \mathcal{O}_X(\ell)) = 0, \quad 0 < i < d, \quad \text{for all } \ell.$$

Remark: A sheaf \mathcal{F} is said to be ACM (arithmetically Cohen-Macaulay) if $\mathcal{F}(\ell)$ has no intermediate cohomology for all ℓ . Over complete intersections, \mathcal{O}_X is ACM.

(iv) Using problem 2, show that the arithmetic genus $p_a(X) = \dim H^d(X, \mathcal{O}_X)$.

4. (Hodge numbers.) For a smooth projective variety over k , set $H^{p,q}(X) = H^q(X, \Omega_X^p)$ and define the Hodge numbers

$$h^{p,q}(X) = \dim_k H^{p,q}(X).$$

Show that for projective space $h^{p,q}(\mathbb{P}^n) = 0$ if $p \neq q$ and $h^{p,q}(\mathbb{P}^n) = 1$ if $p = q$.

Remark: If X is a smooth complex projective variety, then the cohomology of X splits as

$$H^k(X, \mathbb{C}) = \bigoplus_{p+q=k} H^{p,q}(X).$$

This is called the Hodge decomposition. The Hodge numbers $h^{p,q}$ satisfy the symmetries

$$h^{p,q} = h^{q,p} \text{ and } h^{p,q} = h^{d-p, d-q}.$$

The latter symmetry follows from Serre duality. The Betti numbers of X can be calculated from the Hodge numbers

$$b_k(X) = \sum_{p+q=k} h^{p,q}.$$

The Hodge numbers (arranged in a rotated square) form the *Hodge diamond* of X .

Hint: You may wish to start with the Euler sequence

$$0 \rightarrow \Omega_{\mathbb{P}^n} \rightarrow \mathcal{O}_{\mathbb{P}^n}(-1) \otimes \mathbb{C}^{n+1} \rightarrow \mathcal{O} \rightarrow 0.$$

It is important to show that if

$$0 \rightarrow E \rightarrow F \rightarrow \mathcal{O} \rightarrow 0$$

is an exact sequence of vector bundles then you also have exactness of

$$0 \rightarrow \Lambda^p E \rightarrow \Lambda^p F \rightarrow \Lambda^{p-1} E \rightarrow 0.$$

5. (The Picard group.) Let X be a variety. Show that the $\text{Pic}(X)$ can be identified with $H^1(X, \mathcal{O}_X^*)$ where \mathcal{O}_X^* is the sheaf of nowhere-zero regular functions.

Hint: Start with trivializations ψ_i of a line bundle $L|_{U_i} \rightarrow \mathcal{O}_{U_i}$ over open sets U_i , and consider $\psi_{ij} = \psi_i \circ \psi_j^{-1}$ over U_{ij} . Show that ψ_{ij} satisfy the cocycle condition and thus define an element in $H^1(X, \mathcal{O}_X^*)$.

6. (On finiteness of cohomology.) Let $X = \mathbb{A}_k^2 \setminus \{(0,0)\}$. Show that $\dim_k H^1(X, \mathcal{O}_X)$ is infinite.

Hint: Use the Cech cover with two open sets $U = \{x \neq 0\}$ and $V = \{y \neq 0\}$.