

Math 220C - Lecture 17

May 4, 2022

Last time

We defined Riemann surfaces (X, \mathcal{O}_X) as ringed spaces

In particular, we defined

(i) holomorphic functions on $U \subseteq X$.

(ii) holomorphic maps $f: X \rightarrow Y$ of Riemann surfaces.

Recall


\exists open cover $X = \bigcup U_\alpha$, $G_\alpha \subseteq \mathbb{C}$ and

$$\phi_\alpha: (U_\alpha, \mathcal{O}_X|_{U_\alpha}) \longrightarrow (G_\alpha, \mathcal{O}_{G_\alpha})$$

isomorphism of ringed spaces.

In concrete terms Let X Riemann surface. Let $X = \bigcup_{\alpha} U_{\alpha}$

s.t. $(U_{\alpha}, \mathcal{O}_X|_{U_{\alpha}}) \cong (G_{\alpha}, \mathcal{O}_{G_{\alpha}})$ via isomorphism φ_{α} .




Let $U_{\alpha\beta} = U_{\alpha} \cap U_{\beta}$. Note $\varphi_{\beta} \varphi_{\alpha}^{-1}: \varphi_{\alpha}(U_{\alpha\beta}) \rightarrow \varphi_{\beta}(U_{\alpha\beta})$.

must be an isomorphism of ringed spaces.

Thus $\varphi_{\beta} \varphi_{\alpha}^{-1}$ is a biholomorphism between open subsets of \mathbb{C} .

Holomorphic functions

Let X be a Riemann surface, $(U_\alpha, G_\alpha, \varphi_\alpha)$ coordinate charts.



We showed last time that

f holomorphic iff $f \circ \varphi_\alpha^{-1}$ is holomorphic in $\varphi_\alpha(U \cap U_\alpha) \neq \emptyset$.

Remark We can also turn this discussion around.

Let X be a **topological space** (Hausdorff, 2nd countable)

$X = \bigcup_{\alpha} U_{\alpha}$ open cover. Assume we are given

- $\varphi_{\alpha} : U_{\alpha} \rightarrow G_{\alpha}$ homeomorphisms, $G_{\alpha} \subseteq \mathbb{C}$ such that
- $\varphi_{\beta} \varphi_{\alpha}^{-1} : \varphi_{\alpha}(U_{\alpha} \cap U_{\beta}) \rightarrow \varphi_{\beta}(U_{\alpha} \cap U_{\beta})$ biholomorphism

These are called **compatible coordinate charts**

Then X becomes a **Riemann surface**.

Issue Define the sheaf \mathcal{O}_X .


Note U open $\Leftrightarrow U \cap U_{\alpha}$ open $\Leftrightarrow \varphi_{\alpha}(U \cap U_{\alpha})$ open in G_{α} .

Declare $f : U \rightarrow \mathbb{C}$ to be a **section of \mathcal{O}_X** provided.

$f \in \mathcal{O}_X(U) \Leftrightarrow f \circ \varphi_{\alpha}^{-1}$ holomorphic in $\varphi_{\alpha}(U \cap U_{\alpha})$, $\forall \alpha$.

Check \mathcal{O}_X is a sheaf & (X, \mathcal{O}_X) is a Riemann surface.

Meromorphic functions



Definition

f meromorphic in U provided $f \varphi_\alpha^{-1}$ meromorphic in $\varphi_\alpha(U \cap U_\alpha)$

\exists sheaf \mathcal{M} of meromorphic functions

$U \rightsquigarrow \{ \text{meromorphic functions in } U \}$.

Zeros, poles, order

We define the order of a pole or a zero for f to be the order of a pole or a zero for $f \varphi_\alpha^{-1}$ at $\varphi_\alpha(p)$ for $p \in U_\alpha$.

Claim This is independent of choice of α .

Subclaim

Let g be meromorphic in U , $a \in U$. Let $\tau: V \rightarrow U$ be a biholomorphism with $\tau(b) = a$, $b \in V$. Then

g has order m at $a \Rightarrow g \circ \tau$ has order m at b .

We use this for $g = f \varphi_\alpha^{-1}$, $a = \varphi_\alpha(p)$

$\tau = \varphi_\alpha \varphi_\beta^{-1}$, $b = \varphi_\beta(p)$.

$\Rightarrow g \circ \tau = f \varphi_\beta^{-1}$.

The subclaim shows that the order thus defined is independent of the choice of α .

Proof of the Subclaim

WLOG $a = b = 0$, else we can translate.

Write $g(z) = z^m G(z)$, $G(0) \neq 0$.

Since $T(0) = 0$ & $T'(0) \neq 0$ since T is biholomorphism, we

have $T(z) = z S(z)$, $S(0) \neq 0$.

$$\begin{aligned} \text{Note } g \circ T(z) &= T(z)^m G(T(z)) \\ &= z^m S(z)^m G(T(z)). \end{aligned}$$

Since $S(z)^m G(T(z)) \Big|_{z=0} = S(0)^m G(0) \neq 0 \Rightarrow$

\Rightarrow order $g \circ T$ at $z=0$ equals m . as needed.

Remarks Essential singularities are defined similarly.

Aside - Divisors on Riemann surfaces

Definition A **divisor** on a Riemann surface X is a formal sum

$$D = \sum_{p \in X} n_p [p] \text{ with } n_p \in \mathbb{Z} \text{ such that}$$

$S = \{ p \mid n_p \neq 0 \}$ is locally finite.

Examples

i $X = \hat{\mathbb{C}}$, $D = 2[0] + 3[\infty] - 5[1]$ divisor on X

ii D is said to be **effective** if $n_p \geq 0 \forall p \in X$

iii Divisors can be formally **added & subtracted**

$$D = \sum n_p [p], \quad E = \sum m_p [p]$$

$\Rightarrow D \pm E = \sum (n_p \pm m_p) [p]$ is a divisor

iv **restrictions**, $U \subseteq X$ open. If

$$D = \sum_{p \in X} n_p [p] \Rightarrow D|_U = \sum_{p \in U} n_p [p]$$

IV \mathcal{F} sheaf of divisors Div.

$$U \longrightarrow \{ \text{divisors in } U \}$$

VI degree. If X is compact, any divisor is a finite sum.

$$D = \sum n_p [p], \quad n_p \in \mathbb{Z}. \Rightarrow \deg D := \sum_p n_p.$$

Principal divisors If f meromorphic in X , define

$$\square \quad \text{div } f = \sum_{z \in X} \text{ord}(f, z) [z]$$

$$= \sum_{z \text{ zero}} \text{mult}_z(f) [z] - \sum_{p \text{ pole}} \text{mult}_p(f) [p]$$

III Check: $\text{div}(fg) = \text{div } f + \text{div } g.$

Example $X = \hat{\mathbb{C}}$. $f = \frac{\prod_{i=1}^m (z - a_i)}{\prod_{i=1}^n (z - b_i)}$ meromorphic function in $\hat{\mathbb{C}}$
 $a_i, b_i \in \mathbb{C}.$

$$\text{div } f = \sum_{i=1}^m [a_i] - \sum_{i=1}^n [b_i] + (n-m) [\infty]$$

$$\Rightarrow \deg \text{div } f = \sum_{i=1}^m 1 - \sum_{i=1}^n 1 + (n-m) = 0.$$