

Math 220 C - Lecture 3

---

April 1, 2022

Last time

$u: \bar{\Delta} \rightarrow \mathbb{R}$  continuous, harmonic in  $\Delta$ .

$\square$   $u(0) = \frac{1}{2\pi} \int_0^{2\pi} u(e^{is}) ds$  Mean Value Property

$\square$   $u(a) = \frac{1}{2\pi} \int_0^{2\pi} u(\Phi(e^{is})) ds$  MVE + Aut  $\Delta$ .  
to recenter.

where  $\Phi: \Delta \rightarrow \Delta$ ,  $\partial\Delta \rightarrow \partial\Delta$ ,  $z \rightarrow \frac{z+a}{1+\bar{a}z}$

with inverse  $\psi: \Delta \rightarrow \Delta$ ,  $\partial\Delta \rightarrow \partial\Delta$ ,  $z \rightarrow \frac{z-a}{1-\bar{a}z}$

Goal Make formula  $\square$  even more explicit.

## Poisson Kernel

$$P_r(\theta) = \sum_{n=-\infty}^{\infty} r^{|n|} \cos n\theta, \quad 0 \leq r < 1 \quad \text{well defined.}$$

## Three additional Formulas

$$\boxed{a} \quad P_r(\theta) = \operatorname{Re} \frac{1+z}{1-z}, \quad z = r e^{i\theta}$$

$$\frac{1+z}{1-z} = 1 + \frac{2z}{1-z} = 1 + 2z(1+z+z^2+\dots)$$

$$= 1 + 2 \sum_{n=1}^{\infty} z^n = 1 + 2 \sum_{n=1}^{\infty} r^n e^{in\theta}$$

$$= 1 + 2 \sum_{n=1}^{\infty} r^n (\cos n\theta + i \sin n\theta)$$

$$\operatorname{Re} \frac{1+z}{1-z} = 1 + 2 \sum_{n=1}^{\infty} r^n \cos n\theta$$

$$= 1 + \sum_{n=1}^{\infty} r^n (e^{in\theta} + e^{-in\theta})$$

$$= \sum_{n=-\infty}^{\infty} r^{|n|} e^{in\theta} = P_r(\theta)$$

$$\boxed{b} \quad P_r(\theta) = \frac{1 - |z|^2}{|1 - z|^2}$$

$$P_r(\theta) \stackrel{\boxed{a}}{=} \operatorname{Re} \frac{1+z}{1-z} = \operatorname{Re} \frac{(1+z)(1-\bar{z})}{(1-z)(1-\bar{z})}$$

$$= \operatorname{Re} \frac{1 - z\bar{z} + z - \bar{z}}{|1-z|^2}$$

↙ imaginary

$$= \frac{1 - |z|^2}{|1-z|^2}$$

$$\boxed{c} \quad P_r(\theta) = \frac{1 - r^2}{1 - 2r \cos \theta + r^2} \quad \text{very useful.}$$

Indeed use  $\boxed{b}$  for  $z = r e^{i\theta}$ :

$$|1 - z|^2 = (1 - r \cos \theta)^2 + (r \sin \theta)^2$$

$$= 1 + r^2 - 2r \cos \theta \quad \& \quad 1 - |z|^2 = 1 - r^2$$

## Poisson's Formula

$u: \bar{\Delta} \rightarrow \mathbb{R}$  continuous & harmonic in  $\Delta$ ,  $a = r e^{i\theta}$

$$u(a) = \frac{1}{2\pi} \int_0^{2\pi} P_r(\theta - t) u(\tau e^{it}) dt.$$

Proof Recall

$$u(a) = \frac{1}{2\pi} \int_0^{2\pi} u(\Phi(\tau e^{is})) ds$$

Change of variables

$$\tau e^{is} = \Psi(\tau e^{it})$$

Main Claim

$$ds = P_r(\theta - t) dt$$

the Poisson kernel arises  
via change of variables

Assuming this, we obtain

$$\begin{aligned} u(a) &= \frac{1}{2\pi} \int_0^{2\pi} u(\Phi \Psi(\tau e^{it})) \cdot P_r(\theta - t) dt \\ &= \frac{1}{2\pi} \int_0^{2\pi} u(\tau e^{it}) P_r(\theta - t) dt, \text{ as needed.} \end{aligned}$$

## Proof of the Main claim

$$ds = \frac{d(e^{is})}{i e^{is}} = \frac{d\psi(e^{it})}{i \psi(e^{it})} \stackrel{\text{chain rule}}{=} \frac{\psi'(e^{it}) \cdot i e^{it} dt}{i \psi(e^{it})} = \frac{\psi'(z) z}{\psi(z)} dt$$

Recall  $\psi(z) = \frac{z-a}{1-\bar{a}z}$ . Taking logarithmic derivatives

$$z \cdot \frac{\psi'(z)}{\psi(z)} = \frac{z}{z-a} + \frac{\bar{a}z}{1-\bar{a}z}$$

$$= \frac{z}{z-a} - \frac{1}{2} + \frac{1}{2} + \frac{\bar{a}z}{1-\bar{a}z}$$

$$= \frac{1}{2} \cdot \frac{z+a}{z-a} + \frac{1}{2} \cdot \frac{1+\bar{a}z}{1-\bar{a}z} \quad \checkmark \quad 1 = z\bar{z}$$

$$= \frac{1}{2} \cdot \frac{z+a}{z-a} + \frac{1}{2} \cdot \frac{z\bar{z} + \bar{a}z}{z\bar{z} - \bar{a}z}$$

$$= \frac{1}{2} \cdot \frac{z+a}{z-a} + \frac{1}{2} \cdot \frac{z + \bar{a}}{z - \bar{a}}$$

$$= \operatorname{Re} \frac{z+a}{z-a} = \operatorname{Re} \frac{1 + \frac{a}{z}}{1 - \frac{a}{z}} = P_r(\theta-t)$$

using  $\boxed{a}$   $d \frac{a}{z} = \frac{r e^{i\theta}}{e^{it}} = r e^{i(\theta-t)}$



Siméon Poisson

(1781 - 1842)

Students:

Liouville, Carnot, Dirichlet

### Poisson Kernel

$$\begin{aligned} P_r(\theta) &= \sum_{n=-\infty}^{\infty} r^{|n|} e^{in\theta} = \frac{1-r^2}{1-2r\cos\theta+r^2} \\ &= \operatorname{Re} \frac{1+z}{1-z} = \frac{1-|z|^2}{|1-z|^2} \quad \text{for } z = re^{i\theta} \end{aligned}$$

### Poisson integral formula

$u: \bar{\Delta} \rightarrow \mathbb{R}$  continuous, harmonic in  $\Delta$ . Then

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} P_r(\theta-t) u(e^{it}) dt$$

Remark We can dilate & translate to work with any disc  $\Delta(o, R)$ .

Theorem  $u : \bar{\Delta}(o, R) \rightarrow \mathbb{R}$  continuous & harmonic in  $\Delta(o, R)$ .

$$u(a + r e^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} \frac{R^2 - r^2}{R^2 - 2Rr \cos(\theta - t) + r^2} u(a + R e^{it}) dt$$

Proof

$$\tilde{u} : \bar{\Delta} \rightarrow \mathbb{R}, \quad \tilde{u}(z) = u(a + Rz)$$

We apply the previous result to  $\tilde{u}$ . Then

$$u(a + r e^{i\theta}) = \tilde{u}\left(\frac{r}{R} e^{i\theta}\right) = \frac{1}{2\pi} \int_0^{2\pi} \frac{P_{\frac{r}{R}}(\theta - t)}{\frac{r}{R}} \tilde{u}(e^{it}) dt$$

$$= \frac{1}{2\pi} \int_0^{2\pi} \frac{1 - \left(\frac{r}{R}\right)^2}{1 - 2\frac{r}{R} \cos(\theta - t) + \left(\frac{r}{R}\right)^2} u(a + R e^{it}) dt$$

$$= \frac{1}{2\pi} \int_0^{2\pi} \frac{R^2 - r^2}{R^2 - 2Rr \cos(\theta - t) + r^2} u(a + R e^{it}) dt$$



## Two Consequences

[7] Schwarz Integral Formula


[11] Harnack Inequality

## Harnack's Inequality

$u: \bar{\Delta}(a, R) \rightarrow \mathbb{R}$  continuous, harmonic in  $\Delta(a, R)$  &  $u \geq 0$ ,

If  $|z - a| = r \Rightarrow$

$$u(a) \cdot \frac{R-r}{R+r} \leq u(z) \leq u(a) \frac{R+r}{R-r}$$



## Proof

We use  $-1 \leq \cos(\theta - t) \leq 1$ .

The two inequalities are similar. For instance, 2<sup>nd</sup> inequality:

$$u(a + r e^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} \frac{R^2 - r^2}{R^2 - 2Rr \cos(\theta - t) + r^2} u(a + R e^{it}) dt$$

$u \geq 0$

$$\leq \frac{1}{2\pi} \int_0^{2\pi} \frac{R^2 - r^2}{R^2 - 2Rr + r^2} \cdot u(a + R e^{it}) dt$$

$$= \frac{1}{2\pi} \int_0^{2\pi} \frac{(R-r)(R+r)}{(R-r)^2} \cdot u(a + R e^{it}) dt$$

$$= u(a) \frac{R+r}{R-r} \quad \text{using Mean Value Property.}$$



Axel Harnack (1851-1888) was a Baltic-German mathematician.

He proved Harnack's inequality for harmonic functions & Harnack's curve theorem in real algebraic geometry.