


Math 220B - Lecture 8

February 5, 2024

Last time



Argela'-Ascoli  $\mathcal{F}$  family of continuous functions on  $U$

$\mathcal{F}$  normal  $\iff$   $\mathcal{F}$  locally equicontinuous and  
locally (uniformly) bounded.

usually we just say "locally bounded"

---

Today — we give the proof.

All functions today are continuous.

## Notation & Preliminaries

$f: U \rightarrow \mathbb{C}$  continuous,  $K \subseteq U$  compact

$$\|f\|_K = \sup_{z \in K} |f(z)|$$

Note

$$\square \quad \|f+g\|_K \leq \|f\|_K + \|g\|_K$$

$$\square \quad f_n \xrightarrow{K} f \iff \|f_n - f\|_K \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Def  $f_n$  is uniformly Cauchy in  $K$  if

$$\forall \varepsilon > 0 \exists N \forall n, m \geq N, \|f_n - f_m\|_K < \varepsilon.$$

Lemma  $f_n$  converges uniformly in  $K$

$$\iff f_n \text{ uniformly Cauchy in } K.$$

(last time)

## Proof of Arzela-Ascoli

" $\Rightarrow$ " Let  $\mathcal{F}$  be normal.

(1)  $\mathcal{F}$  locally bounded

Let  $K \subseteq U$  compact. We show  $\mathcal{F}|_K$  bounded. i.e.

$$\exists M > 0 \quad \forall f \in \mathcal{F} \Rightarrow \|f\|_K < M.$$

Assume not for a contradiction. Then

$$\forall M > 0 \quad \exists f_m \in \mathcal{F} \text{ with } \|f_m\|_K \geq M$$

Letting  $M = n$ , we obtain a sequence  $f_n$  with  $\|f_n\|_K \geq n$ .

Since  $\mathcal{F}$  normal, we can find a subsequence  $f_{n_k} \xrightarrow{K} f$

Thus  $\|f_{n_k} - f\|_K < 1$  if  $k$  sufficiently large.

Note  $f_{n_k}$  continuous  $\Rightarrow f$  continuous. so  $\|f\|_K < \infty$ . Then

$$\|f\|_K \geq \|f_{n_k}\|_K - \|f_{n_k} - f\|_K \geq n_k - 1 \rightarrow \infty \text{ as } k \rightarrow \infty$$

This gives a contradiction.

## (2) $\mathcal{F}$ locally equicontinuous

Let  $K \subseteq U$  compact. We show  $\mathcal{F}|_K$  uniformly equicontinuous.

that is  $\forall \varepsilon \exists \delta : \forall x, y \in K, |x - y| < \delta \ \forall f \in \mathcal{F}$  then

$$|f(x) - f(y)| < \varepsilon.$$

Assume not, then

$\exists \varepsilon \ \forall \delta \exists x_\delta, y_\delta \in K$  with  $|x_\delta - y_\delta| < \delta \ \exists f_\delta \in \mathcal{F}$  but

$$|f_\delta(x_\delta) - f_\delta(y_\delta)| \geq \varepsilon.$$

Take  $\delta = \frac{1}{n}$ . Then

$\exists x_n, y_n \in K, |x_n - y_n| < \frac{1}{n} \ \exists f_n \in \mathcal{F}$  with


$$|f_n(x_n) - f_n(y_n)| \geq \varepsilon.$$

After passing to a subsequence & relabelling, we arrange

□  $f_n \xrightarrow{K} f$  because  $\mathcal{F}$  normal

□  $|x_n - y_n| < \frac{1}{n}$

□  $|f_n(x_n) - f_n(y_n)| \geq \varepsilon.$



Using  $f_n$  continuous,  $f_n \Rightarrow f$  we get  $f$  continuous.

Since  $K$  compact  $\Rightarrow f|_K$  uniformly continuous.

Then  $\exists \tau > 0$  with

$$|x - y| < \tau, \quad x, y \in K \Rightarrow |f(x) - f(y)| < \frac{\epsilon}{3}. \quad (1)$$

Let  $N$  be so that  $\forall n \geq N$ , we have  $\frac{1}{n} < \tau$  and

$$\|f_n - f\|_K < \frac{\epsilon}{3}. \quad (2)$$

Then  $|x_n - y_n| < \frac{1}{n} < \tau \Rightarrow |f(x_n) - f(y_n)| < \frac{\epsilon}{3}$  by (1).

$|f_n(x_n) - f(x_n)| < \frac{\epsilon}{3}$  &  $|f_n(y_n) - f(y_n)| < \frac{\epsilon}{3}$  by (2).

By triangle inequality (see picture)

$$|f_n(x_n) - f_n(y_n)| < \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon$$

contradicting  $\square$

## The Converse

Assume  $\mathcal{F}$  is locally equicontinuous & locally bounded.

$\stackrel{?}{\implies} \mathcal{F}$  normal

---

Let  $f_n \in \mathcal{F}$ . We wish to find a subsequence converging locally uniformly?

How do we find such a subsequence?

---

Plan [1] arrange pointwise convergence of  $f_n$

[2] show local uniform convergence using local equicontinuity

Better Plan [1] arrange pointwise convergence of  $f_n$  only

at a countable dense set

[2] show local uniform convergence

Let  $\{a_k\}$  be the set of points in  $U$  with rational coordinates enumerated in any order. **Dense!**

---

Claim [L] After passing to a subsequence of  $f_n$  & relabelling, we may assume

(\*)  $\forall k$ , the sequence  $f_n(a_k)$  converges as  $n \rightarrow \infty$ .

---

Claim [L] If  $\{f_n\}$  locally equicontinuous & (\*) holds

$\Rightarrow f_n$  converges locally uniformly.

We win!



## Proof of Claim 14 Cantor diagonalization

We only use pointwise boundedness of  $\{f_n\}$ .

---

Consider  $f_1(a_1), f_2(a_1), \dots, f_n(a_1), \dots$  bounded

Find a subsequence

$(s_1)$   $f_{11}, f_{12}, \dots, f_{1n}, \dots$   $\swarrow$  converges at  $a_1$


Look at the values of  $(s_1)$  at  $a_2$  & repeat. We find

$(s_2)$   $f_{21}, f_{22}, \dots, f_{2n}, \dots$   $\swarrow$  converges at  $a_2$   
and  $a_1$

Look at the values of  $(s_2)$  at  $a_3$  & repeat.

---

We obtain an array:




Each row is a subsequence of the previous one.

Consider the diagonal subsequence

$$f_{11}, f_{22}, f_{33}, \dots, f_{nn}, \dots$$

It is a subsequence of the original sequence, &

converges at each  $a_k$ . Indeed



## Proof of Claim [ii]

Know [a]  $\{a_k\}$  dense in  $U$  and

$\forall k$ , the sequence  $\{f_n(a_k)\}_n$  converges

[b]  $f_n$  locally equicontinuous

Wish  $\forall \alpha \in U$ ,  $\exists \Delta =$  bounded open ball in  $U$ ,  $\alpha \in \Delta$

$f_n|_{\Delta}$  converges uniformly.

(1)  $\forall \alpha \exists \Delta \ni \alpha \in \bar{\Delta}$ ,  $\mathcal{F}|_{\Delta}$  equicontinuous. (uniformly)

Thus  $\forall \varepsilon \exists \delta: \forall |x-y| < \delta, x, y \in \bar{\Delta}, \forall f \in \mathcal{F}$


$$|f(x) - f(y)| < \varepsilon/3$$

(2)  $\bar{\Delta}$  can be covered by  $\Delta_i = \Delta(a_i, \delta)$  for  $a_i \in \bar{\Delta}$ .

This because  $\{a_i\} \cap \bar{\Delta}$  is dense in  $\bar{\Delta}$ .

By compactness, we may assume

$$\bar{\Delta} \subseteq \bigcup_{i=1}^e \Delta(a_i, \delta).$$



(3) Since  $\{f_n(a_i)\}_{i=1,2}$  is convergent, it is Cauchy. Hence

$$\forall \varepsilon \exists N \forall n, m \geq N \forall 1 \leq i \leq l$$

$$|f_n(a_i) - f_m(a_i)| < \varepsilon/3$$

(4) Let  $z \in \bar{\Delta}$ . By (2),  $\exists i$  with  $|z - a_i| < \delta$ . Let  $n, m \geq N$ .

as in (3). Then

$$\begin{aligned} |f_n(z) - f_m(z)| &\leq \overset{\text{use (1)}}{|f_n(z) - f_n(a_i)|} + \overset{\text{use (3)}}{|f_n(a_i) - f_m(a_i)|} + \overset{\text{use (1)}}{|f_m(a_i) - f_m(z)|} \\ &< \varepsilon/3 + \varepsilon/3 + \varepsilon/3 \end{aligned}$$

(5) Conclusion  $\|f_n - f_m\|_{\bar{\Delta}} < \varepsilon \forall n, m \geq N$ .

$\Rightarrow \{f_n\}$  uniformly Cauchy in  $\bar{\Delta}$

Lemma  $\Rightarrow \{f_n\}$  converges uniformly in  $\bar{\Delta}$ .

This completes the proof.

Remark The converse only used pointwise boundedness

$\mathcal{F}$  normal  $\Leftrightarrow \mathcal{F}$  pointwise bounded + locally equicont

$\Leftrightarrow \mathcal{F}$  locally bounded + locally equicont.

The second version bears connections with Montel & it is more uniform.

End of Part I of the course