

Definition (of base change): Given $X \xrightarrow{\pi_X} S$

morphism of schemes, the base change of X to a scheme $S' \xrightarrow{\varphi} S$ is the fiber product

$$X' := X \times_S S' \xrightarrow{\quad} X$$
$$\downarrow \quad \square \quad \downarrow \pi_X$$
$$S' \xrightarrow{\varphi} S$$

Definition (of (universally) closed morphisms):

A morphism of schemes is closed if the image of any closed subset is closed. A morphism $f: X \rightarrow Y$ of schemes is universally closed if for any $Y' \xrightarrow{g} Y$, the base change $X \times_Y Y' \xrightarrow{f'} Y'$ is closed.

Example: $A' \rightarrow \text{Spec } k$ is closed, but it is
not universally closed: base change to $A'^2 \rightarrow \text{Spec } k$:

$$A'^2 \cong A' \times_{\bar{k}} A' \xrightarrow{\pi_1} A'$$

$$\pi_2 \downarrow \quad \square \quad \downarrow$$

$$A' \longrightarrow \text{Spec } k$$

π_2 is not closed: the
image of the hyperbola
 $z(xy-1) \subset A' \times A'$
is not closed in A' .

Def: A morphism is proper if it is of finite type,
separated and universally closed.

Theorem: The valuative criterion of properness:

Let $f: X \rightarrow Y$ be a morphism of finite type,
with X noetherian. Then f is proper if and only if,

for any field K and valuation ring $R \subset K$,
 and any morphisms $T := \text{Spec } R \rightarrow Y$, $U := \text{Spec } k \rightarrow X$
 forming a commutative diagram

$$\begin{array}{ccc} U & \longrightarrow & X \\ \downarrow & \nearrow f! & \downarrow f \\ T & \longrightarrow & Y \end{array}$$

there is a unique morphism $T \rightarrow X$ making the
 whole diagram above commutative.

Proof: First assume $f: X \rightarrow Y$ is proper.

For any data as in the theorem

since f is by definition separated,

\exists at most one lift $T \rightarrow X$ making the diagram commute

$$\begin{array}{ccc} U & \longrightarrow & X \\ \downarrow & \nearrow f! & \downarrow f \\ T & \longrightarrow & Y \end{array}$$

So we need to prove the existence of the lift.

$$\text{Spec } k = U \longrightarrow X$$

\downarrow $\dashv \pi_1^*$ $\downarrow f$

$$\text{Spec } R = T \longrightarrow Y$$

Let us do the base change

$$\begin{array}{ccc} X \times_T Y & \xrightarrow{\alpha_1} & X \\ \pi_1 \downarrow & & \downarrow f \\ T & \xrightarrow{\quad} & Y \end{array}$$

$$\begin{array}{ccc} U & \xrightarrow{\quad} & X \times_T Y \xrightarrow{\alpha_1} X \\ \downarrow & \nearrow & \downarrow \pi_1 \\ & X \times_T Y & \xrightarrow{\alpha_1} X \\ \pi_2 \downarrow & & \downarrow f \\ T & \xrightarrow{\quad} & Y \end{array}$$

$U \rightarrow X \times_T Y$ exists and is unique by the universal property of fiber products.

Let $\xi_1 \in X \times_T Y$ be the image of \cup , let
 $Z := \overline{\{\xi_1\}}$ be the closure of ξ_1 in $X \times_Y T$ with the
reduced induced scheme structure.

If universally closed $\Rightarrow \mu_2$ is closed $\Rightarrow \mu_2(Z) \subset T$
is closed

$\mu_2(Z)$ contains $\mu_2(\xi_1)$ which is the generic point of T

$$\Rightarrow \mu_2(Z) = T$$

$\Rightarrow \exists \xi_0 \in Z$ s.t. $\mu_2(\xi_0)$ = the closed point of T .

$\mu_2 : Z \rightarrow T$ \Rightarrow local hom. of local rings
 $\xi_1 \mapsto t_1$
 $\xi_0 \mapsto t_0$

$$R = \mathcal{O}_{T, t_0} \rightarrow \mathcal{O}_{Z, \xi_0}$$

$$K = \mathcal{O}_{T, \xi_1} \rightarrow \mathcal{O}_{Z, \xi_1} = K$$

$\Rightarrow G_{Z, \xi_0}$ dominates R

$\Rightarrow G_{Z, \xi_0} = R$ because R is a valuation ring.
(maximal for dominance)

By the lemma (4.4 in Hartshorne) from one or two lectures ago,

\exists morphism $T \xrightarrow{\quad} X \times_T Y$ sending t_0 to \mathfrak{f}_0 and t_1 to \mathfrak{f}_1 .

Now compose with π_1 to obtain the desired lift

$T \xrightarrow{\quad} X$.

Conversely, suppose f is of finite type, X noetherian and the valuative criterion holds.

By the valuative criterion of separatedness, we know f is separated.

We need to prove that f is universally closed.

For any $\gamma' \rightarrow Y$, we show that $X \times_{Y'} \gamma' \xrightarrow{p_2} \gamma'$ is closed.

$$X' := X \times_{Y'} \gamma' \xrightarrow{p_1} X$$

$$f' := p_2 \downarrow \square \downarrow f \\ \gamma' \longrightarrow Y$$

Let $Z \subset X'$ be a closed subset. $f'(Z) \subset Y'$
Endow Z with the reduced induced scheme structure.

Exercise: f' is of finite type, and $f'|_Z$ is of finite type.

$\Rightarrow f'|_Z$ is quasi-compact.

We use a lemma (4.5 in Hartshorne) from previous lectures:
 $f'(Z)$ is closed iff it is closed under specialization.

Let $\beta_1 \in Z$, put $y_1 := f(\beta_1)$

For any $y_0 \in \overline{\{y\}}$ a specialization, we show $y_0 \in f'(Z)$.

Endow $W := \{y_1\}$ with its reduced induced scheme structure.

$$\mathcal{O}_{W, y_0} \hookrightarrow \mathcal{O}_{W, y_1} = K_W \hookrightarrow K := k(\beta_1)$$

residue field in z

Let R be a valuation ring of K dominating \mathcal{O}_{W, y_0} .

Apply Lemma 4.4 again to obtain a morphism

$$\begin{array}{ccccc} U = \text{Spec } K & \hookrightarrow & T = \text{Spec } R & \xrightarrow{\quad} & Y' \\ & \swarrow & \uparrow t_1 & \mapsto & y_1 = f(\beta_1) \\ & & t_0 & \mapsto & y_0 \end{array}$$

By construction, we have the commutative diagram

$$\begin{array}{ccccc} t_1 & \longrightarrow & \mathcal{Z} & \ni & \mathcal{Z} \\ U = \text{Spec } K & \longrightarrow & Z & \ni & \mathcal{Z} \\ \downarrow & \text{?} & \downarrow f'|_Z & & \\ T = \text{Spec } R & \longrightarrow & Y' & & \\ t_0 & \longleftarrow & \mathcal{Y}' & & \\ t_1 & \longleftarrow & \mathcal{Y} & & \end{array}$$

Compose: $t_i \in U \longrightarrow Z \hookrightarrow X' \longrightarrow X$

$$\begin{array}{ccccc} & & X' & \longrightarrow & X \\ & & \downarrow f' & \nearrow f'|_Z & \downarrow f \\ t_0, t_1 \in T & \longrightarrow & Y' & \longrightarrow & Y \\ & & \downarrow & & \downarrow f \\ & & \mathcal{Y} & & \end{array}$$

By the valuative criterion, $\exists!$ lift $T \xrightarrow{\quad} X$
making the diagram commutative. $\Rightarrow T \xrightarrow{\quad} X'$ by the
universal property of fiber products.

The generic point of T maps into Z , and Z is closed, hence all of T maps into Z .

Let $z_0 \in Z$ be the image of $t_0 \in T$.

Then $f'(z_0) = \text{image of } t_0 = y_0 \in f'(Z)$.

□