

Recall: Theorem: X irreducible, separated, of finite type over an algebraically closed field k . Then $\Omega'_{X/k}$ is a locally free sheaf of rank = $\dim X$ iff X is a nonsingular variety / k .

Reason: $\forall x$ closed $\Omega'_{X/k, x} \otimes_{\mathcal{O}_{X, x}} k \cong \mathfrak{m}_x / \mathfrak{m}_x^2$

Recall: If $Y \subset X$ is a closed subscheme, then we have

$$\begin{array}{ccc} Y & \subset & X \\ \downarrow & & \downarrow \\ S & & S \end{array}$$

the natural exact sequence

$$\mathcal{I}/\mathcal{I}^2 \longrightarrow \Omega'_{X/S} \Big|_Y \longrightarrow \Omega'_{Y/S} \longrightarrow 0$$

pull-back via inclusion

Theorem (8.17) If $S = \text{Spec } k$ k alg. closed and X is a nonsingular variety, then Y is nonsingular

iff

(1) $\Omega_{Y/k}^1$ is locally free

(2) the natural sequence

$$0 \rightarrow \mathcal{I}/\mathcal{I}^2 \rightarrow \Omega_{X/k}^1|_Y \rightarrow \Omega_{Y/k}^1 \rightarrow 0$$

is exact.

Furthermore, if the above holds, then \mathcal{I} is locally generated by $d = \text{codim}_X(Y)$ elements and $\mathcal{I}/\mathcal{I}^2$ is locally free of rank d .

X variety \mathbb{k}

Def: The tangent sheaf of X is

$$\mathcal{T}_X := \mathcal{H}om_{\mathcal{O}_X}(\Omega_X^1/\mathbb{k}, \mathcal{O}_X)$$

note: $\forall x \in X$ closed point, $\mathcal{T}_{X,x} \otimes_{\mathcal{O}_{X,x}} \mathbb{k} \cong (\mathcal{M}_x/\mathcal{M}_x^2)^* = T_x X$

If $Y \subset X$ is a closed subscheme, the normal sheaf of Y in X is $\mathcal{N}_{Y/X} := (\mathcal{I}/\mathcal{I}^2)^* := \mathcal{H}om_{\mathcal{O}_X}(\mathcal{I}/\mathcal{I}^2, \mathcal{O}_X)$

The sheaf $\mathcal{I}/\mathcal{I}^2$ is called the conormal sheaf.


If Y is also a nonsingular variety, then we can dualize the exact sequence

$$0 \rightarrow \mathcal{I}/\mathcal{I}^2 \rightarrow \Omega'_X|_Y \rightarrow \Omega'_Y \rightarrow 0 \quad /k$$

to obtain

$$0 \rightarrow \mathcal{C}_Y \rightarrow \mathcal{C}_X|_Y \rightarrow \mathcal{N}'_{Y/X} \rightarrow 0$$

at closed points: $0 \rightarrow T_x Y \rightarrow T_x X \rightarrow N_{Y/X, x} \rightarrow 0$



Def: If $Y \subset X$ is a closed subscheme of a nonsingular variety, then we say Y is a local complete intersection in X if the ideal sheaf \mathcal{I}_Y can be locally generated by $d = \text{codim}_X(Y)$ elements.

Prop.: If $Y \subset X$ is a local complete intersection, (l.c.i.) then $\mathcal{I}_Y/\mathcal{I}_Y^2$ is locally free of rank d

and the sequence

$$0 \rightarrow \mathcal{I}_Y/\mathcal{I}_Y^2 \rightarrow \Omega_X^1|_Y \rightarrow \Omega_Y^1 \rightarrow 0$$

is exact.

$$\left(\Rightarrow \wedge^n \Omega_X^1|_Y \cong \wedge^{n-d} \Omega_Y^1 \otimes \wedge^d (\mathcal{I}_Y/\mathcal{I}_Y^2) \right)$$

Def: X nonsingular variety / k

The canonical sheaf of X is

$$K_X := \Lambda^n \Omega^1_X \quad \text{where } n = \dim X.$$

Note: When $Y \subset X$ is nonsingular, then

$$K_Y \cong K_X \otimes \Lambda^d \mathcal{N}_{Y/X}$$

Note: Being l.c.i. is independent of the ambient scheme X .

Cohen-Macaulay is a generalization of l.c.i.

Let A be a ring, M an A -module

Def: A sequence a_1, \dots, a_n of elements of A is called M -regular if a_1 is not a zero divisor in M , i.e., multiplication by a_1 : $M \xrightarrow{a_1} M$ is injective, a_2 is not a zero divisor in M/a_1M , a_3 is not a zero divisor in $M/a_1M + a_2M$, \dots

If A is a local ring with maximal ideal \mathfrak{m} , the depth of M is the maximum length of a regular sequence of elements of \mathfrak{m} .

We say that A is a Cohen-Macaulay ring if

$$\text{depth } A = \dim A.$$

Note: We always have $\text{depth} \leq \dim$.

- Facts: (1) Regular local rings are Cohen-Macaulay, quotients of Cohen-Macaulay rings by regular sequences are Cohen-Macaulay. In particular, local rings of l.c.i. schemes are Cohen-Macaulay.
- (2) Noetherian local rings of $\dim \leq 1$ (i.e.; ^{local} Artinian rings) are Cohen-Macaulay.
- (3) One-dimensional reduced noetherian local rings are Cohen-Macaulay.
- (4) Two-dimensional integrally closed noetherian local domains are Cohen-Macaulay.

(5) If A is a finitely generated Cohen-Macaulay algebra over a field k with an action of a finite group G , then the subring $A^G \hookrightarrow A$ of G -invariants is Cohen-Macaulay.

(6) Determinantal rings are Cohen-Macaulay:

For any ring A , given an $n \times n$ matrix $(a_{ij}) =: M$ with entries in A , consider the ideal I generated by the minors of size $r \times r$ of M : this is a determinantal ideal. A ring is called determinantal if it is the quotient of a regular local ring by a determinantal ideal.

Theorem (8.22A Serre):

A noetherian ring A is normal (every localization of A at a prime ideal is an integrally closed domain)

iff it satisfies

- $R_1 \rightarrow$ (1) \forall prime $\mathfrak{p} \subset A$ of height ≤ 1 , $A_{\mathfrak{p}}$ is regular
- $S_2 \rightarrow$ (2) \forall prime $\mathfrak{p} \subset A$ of height ≥ 2 , $\text{depth}(A_{\mathfrak{p}}) \geq 2$.

\Rightarrow Prop. (8.23) If $Y \subset X$ is a l.c.i. / k . Then

(1) Y is Cohen-Macaulay

(2) Y is normal iff it is regular in codim. 1. (R_1)

If a scheme is Cohen-Macaulay, S_2 is automatically satisfied.