

So we have commutative diagrams with exact rows:

$$\begin{array}{ccccccc}
 0 \rightarrow & \check{H}^0(\mathcal{U}, \mathcal{F}) & \rightarrow & \check{H}^0(\mathcal{U}, \mathcal{G}) & \rightarrow & \check{H}^0(\mathcal{U}, \mathcal{R}) & \rightarrow & \check{H}^1(\mathcal{U}, \mathcal{F}) & \rightarrow & 0 \\
 & \cong \downarrow \kappa & & \cong \downarrow \kappa & & \cong \downarrow \kappa & & \downarrow \kappa & & \\
 0 \rightarrow & H^0(X, \mathcal{F}) & \rightarrow & H^0(X, \mathcal{G}) & \rightarrow & H^0(X, \mathcal{R}) & \rightarrow & H^1(X, \mathcal{F}) & \rightarrow & 0
 \end{array}$$

(note  $\check{H}^i(\mathcal{U}, \mathcal{G}) = H^i(X, \mathcal{G}) = 0 \quad \forall i > 0$  because  $\mathcal{G}$  is flasque)

$$\Rightarrow \kappa: \check{H}^1(\mathcal{U}, \mathcal{F}) \xrightarrow{\cong} H^1(X, \mathcal{F})$$

For  $i \geq 1$ , we have commutative diagrams:

$$\begin{array}{ccccccc}
 0 \rightarrow & \check{H}^i(\mathcal{U}, \mathcal{R}) & \xrightarrow{\delta} & \check{H}^{i+1}(\mathcal{U}, \mathcal{F}) & \rightarrow & 0 \\
 & \downarrow \kappa & & \downarrow \kappa & & \\
 0 \rightarrow & H^i(X, \mathcal{R}) & \rightarrow & H^{i+1}(X, \mathcal{F}) & \rightarrow & 0
 \end{array}$$

We know  $\kappa: \check{H}^i(\mathcal{U}, \mathcal{F}) \xrightarrow{\cong} H^i(X, \mathcal{F}) \quad \forall$   
quasi-coherent sheaves on  $X$ . By induction, we  
obtain  $\kappa: \check{H}^i(\mathcal{U}, \mathcal{F}) \xrightarrow{\cong} H^i(X, \mathcal{F}) \quad \forall i$   
and all quasi-coherent sheaves.  $\square$ .

Benchmark: When  $X$  is a separated Noetherian scheme,  
it satisfies the hypothesis of the theorem for any  
open affine covering by Ex. II.4.3 from last quarter's homework.

Corollary: If  $X$  has a covering by  $r+1$  open affine  
sets s.t. all the intersections are affine, then  
 $\forall$  quasi-coherent sheaves  $\mathcal{F}$  and all  $i > r$ ,  
 $H^i(X, \mathcal{F}) = 0$ .

We use the previous to compute cohomology on projective space:

Let  $A$  be a noetherian ring.

$$S := A[X_0, \dots, X_n]$$

$$X := \mathbb{P}_A^n = \text{Proj } S$$

$\forall \mathcal{F}$  sheaf of  $\mathcal{O}_X$ -modules:  $\Gamma_*(\mathcal{F}) = \bigoplus_{n \in \mathbb{Z}} H^0(X, \mathcal{F}(n))$

Recall  $\Gamma_*(\mathcal{O}_X) = S$  as graded  $S$ -modules.

and  $\exists$  natural map

$$\widetilde{\Gamma_*(\mathcal{F})} \longrightarrow \mathcal{F}$$

which is an isomorphism if  $\mathcal{F}$  is quasi-coherent.

Theorem: (1)  $\forall n \in \mathbb{Z}, \forall i \neq 0, n, H^i(X, \mathcal{O}_X(n)) = 0$   
(2)  $H^n(X, \mathcal{O}_X(-n-1)) \cong A$   
(3)  $\forall n \in \mathbb{Z}$ , the natural map (cup-product)

$$H^0(X, \mathcal{O}_X(n)) \times H^n(X, \mathcal{O}_X(-n-1)) \rightarrow H^n(X, \mathcal{O}_X(-n-1)) = A$$

is a perfect pairing of finitely generated free  $A$ -modules.

Proof: Put  $\mathcal{F} := \bigoplus \mathcal{O}_X(n)$ . Then  $\mathcal{F}$  is quasi-coherent, in fact, locally free of infinite rank.  $\mathcal{F}$  has a  $\mathbb{Z}$ -grading.

We compute  $H^i(\mathcal{F})$ , keeping track of the  $\mathbb{Z}$ -grading.

(cohomology commutes with arbitrary direct sums: they are direct limits of finite direct sums).

Recall that all cohomology groups are

$$H^0(X, \mathcal{O}_X) = A \text{ - modules.}$$

We compute  $\check{H}^i(\mathcal{U}, \mathcal{F})$  for  $\mathcal{U} = \{U_0, \dots, U_n\}$  the usual open cover of  $X$ .

Recall  $U_i = D_+(X_i) \subset X$

$$U_{i_0, \dots, i_r} = D_+(X_{i_0} X_{i_1} \dots X_{i_r}) \subset X$$

$\forall \{i_0, \dots, i_r\}$ , we have

$$\mathcal{F}(U_{i_0, \dots, i_r}) = \bigoplus_{n \in \mathbb{Z}} \mathcal{O}_X(n)(U_{i_0, \dots, i_r})$$

$$= S[X_{i_0} \dots X_{i_r}]^{-1}$$

because  $\mathcal{O}_X(n)(U_{i_0, \dots, i_r}) = S[X_{i_0} \dots X_{i_r}]^{-1}_{\text{degree } n}$

So we have the Čech complex of  $\mathcal{F}$ :

$$0 \rightarrow \prod_{i=0}^n \mathcal{F}(U_i) \rightarrow \prod_{0 \leq i < j \leq n} \mathcal{F}(U_{ij}) \rightarrow \dots \rightarrow \mathcal{F}(U_{0 \dots n}) \rightarrow 0$$

$\parallel$   $\parallel$   $\parallel$

$$0 \rightarrow \prod_{i=0}^n S[x_i^{-1}] \rightarrow \prod_{0 \leq i < j \leq n} S[x_i^{-1} x_j^{-1}] \rightarrow \dots \rightarrow S[x_0^{-1} \dots x_n^{-1}] \rightarrow 0$$

The kernel of the first map is  $H^0(X, \mathcal{F}) = S$  (already known).  $H^n(X, \mathcal{F})$  is the cokernel of the last map:

$$\prod_{i=0}^n S[x_0^{-1} \dots \widehat{x_i^{-1}} \dots x_n^{-1}] \rightarrow S[x_0^{-1} \dots x_n^{-1}]$$

The ring  $S[x_0^{-1} \dots x_n^{-1}]$  is the free  $A$ -module with basis all the Laurent monomials, i.e., monomials with both

positive and negative powers.

The cokernel of the map is the free  $A$ -module with basis the monomials where all the  $X_i$  have negative power:  $\{X_0^{l_0} \cdots X_n^{l_n}, l_i < 0 \forall i\}$

The grading on  $\mathcal{F}$  or  $S[X_0^{-1} \cdots X_n^{-1}]$  is given by  $\sum_{i=0}^n l_i$ . So  $H^n(X, \mathcal{O}_X(-n)) = 0$  if  $n \leq n$

and  $H^1(X, \mathcal{O}_X(-n-1-n))$  is isomorphic to the

free  $A$ -module with basis  $X_0^{l_0} \cdots X_n^{l_n}$   $l_i < 0$   
 $\sum_{i=0}^n l_i = -n-1-n$   
 $\sum_{i=0}^n l_i = -n-1-n$

The pairing between  $H^0(X, \mathcal{O}_X(n))$  and

$H^n(X, \mathcal{O}_X(-n-1-n))$  is induced by multiplication:

$$\left( \bigoplus_{\substack{l_i \geq 0 \\ \sum l_i = n}} A X_0^{l_0} \cdots X_n^{l_n} \right) \times \left( \bigoplus_{\substack{m_i < 0 \\ \sum m_i = -n-1-n}} A X_0^{m_0} \cdots X_n^{m_n} \right)$$

$\xrightarrow{\hspace{15em}} A$   
 the dual of  $X_0^{l_0} \cdots X_n^{l_n}$  is  $X_0^{-l_0-1} \cdots X_n^{-l_n-1}$ .

This defines a perfect pairing.

It remains to prove that  $H^i(X, \mathcal{F}) = 0$  for  $0 < i < n$ .

Localize the Čech complex at  $X_n$ : this produces an exact sequence <sup>except on the left</sup>, because the cohomology of the localization at  $X_n$  is the cohomology of  $\mathcal{F}|_{U_n}$  which is 0 because  $U_n$  is affine and  $\mathcal{F}$  is quasi-coherent.

→ in positive degrees.



$$\Rightarrow \forall i > 0 \quad H^i(X, \mathcal{O}_X)[X_n^{-1}] = 0$$

(localization is exact).

We prove, by induction on  $n$ , that multiplication by  $X_n$  is injective on  $H^i(X, \mathcal{O}_X)$ . This will imply

that  $H^i(X, \mathcal{O}_X) = 0$  for  $0 < i < n$ .

$$Z(X_n) \cong \mathbb{P}^{n-1} \hookrightarrow \mathbb{P}^n$$

$$\text{recall } \mathcal{I}_{Z(X_n)} \cong \mathcal{O}_X(-Z(X_n)) \cong \mathcal{O}_X(-1)$$

$\Rightarrow$  exact sequence

$$0 \longrightarrow \mathcal{I}_{Z(X_n)} \xrightarrow{X_n} \mathcal{O}_X \longrightarrow \mathcal{O}_{Z(X_n)} \longrightarrow 0$$

$$0 \longrightarrow \mathcal{O}_X(-1) \xrightarrow{X_n} \mathcal{O}_X \longrightarrow \mathcal{O}_Z \longrightarrow 0$$

$$Z = Z(X_n)$$

$$\Rightarrow \quad \forall n$$

$$0 \rightarrow \mathcal{O}_X(-n-1) \xrightarrow{X_n} \mathcal{O}_X(n) \rightarrow \mathcal{O}_Z(n) \rightarrow 0$$

induction hypothesis:  $H^i(Z, \mathcal{O}_Z(n)) = 0 \quad \forall n$   
 $\forall 0 < i < n-1$

Furthermore, the map

$$H^0(X, \mathcal{O}_X(n)) \rightarrow H^0(Z, \mathcal{O}_X(n))$$

is surjective hom. pol.

in  $X_0, \dots, X_n$  of degree  $n$

hom. pol. in  
 $X_0, \dots, X_{n-1}$  of degree  $n$ .

$$\Rightarrow H^1(X, \mathcal{O}_X(n-1)) \xrightarrow{X_n} H^1(X, \mathcal{O}_X(n)) \text{ is injective}$$

and  $H^i(X, \mathcal{O}_X(n-1)) \xrightarrow{X_n} H^i(X, \mathcal{O}_X(n))$  is also

injective  $\forall 1 < i < n$  because  $H^{i-1}(Z, \mathcal{O}_Z(n)) = 0$   
 by induction. □