Math 20E

August 13, 2014

**Question 1** In order for a transformation  $T: R \to S$  to be a coordinate transformation that can be used to change variables in a double or triple integral, it should

- **A.** be a one-to-one mapping mapping of R
- **B.** map R onto S
- C. both A and B
- \*D. both  $\bf A$  and  $\bf B$ , except that would be OK if it failed to be one-to-one on parts of the boundary of R
- **E.** none of the above: "one-to-one" and "onto" have nothing to do with coordinate transformations.

**Question 2** Given domains  $D \subset \mathbb{R}^2$  and  $S \subset \mathbb{R}^2$  and a one-to-one transformation  $T:D \to S$  that maps D onto S. Then T can be used to change variables as follows:

**A.** 
$$\iint_{S} f(x,y) dx dy = \iint_{D} f(T(u,v)) |\det[\mathbf{D}T(u,v)]| du dv.$$

**B.** 
$$\iint_D f(u,v) du dv = \iint_S f(T(x,y)) |\det[\mathbf{D}T(x,y)]| dx dy.$$

C. 
$$\iint_D f(u,v) du dv = \iint_S f\left(T^{-1}(x,y)\right) \left| \det \left[ \mathbf{D} T^{-1}(x,y) \right] \right| dx dy.$$

D. Both A and B

\*E. Both A and C

 ${\bf Question~3}~$  The speed of an object is constant. The object's

- \*A. velocity and acceleration are perpendicular.
- **B.** acceleration is zero.
- C. velocity is constant.
- D. both B and C.