

Math 20E

August 20, 2014

Question 1 The surface $x^2 + y^2 + z = 1$ for $z \geq 0$ is parametrized by $\Phi : D \rightarrow \mathbb{R}^3$, where D is the unit disk $u^2 + v^2 \leq 1$ and $\Phi(u, v) = (u, v, 1 - u^2 - v^2)$. Then, $\mathbf{T}_u \times \mathbf{T}_v = (2u, 2v, 1)$ and

- A.** Φ is a one-to-one mapping of D onto $S = \Phi(D)$.
- B.** The parametrized surface Φ is regular at every point of S .
- C.** The surface S has a tangent plane at every point.
- *D.** **A, B** and **C**
- E.** none of the above

Question 2 The surface $x^2 + y^2 + z = 1$ for $z \geq 0$ is parametrized by $\Psi : R \rightarrow \mathbb{R}^3$, where R is the rectangle $[0, 1] \times [0, 2\pi]$ and $\Psi(u, v) = (u \cos(v), u \sin(v), 1 - u^2)$. Then, $\mathbf{T}_u \times \mathbf{T}_v = u(2u \cos(v), 2u \sin(v), u)$ and

- A.** Ψ is a one-to-one mapping of R onto $S = \Psi(R)$.
- B.** The parametrized surface Ψ is regular at every point of S .
- ***C.** The surface S has a tangent plane at every point.
- D.** **A, B** and **C**
- E.** none of the above

Question 3 The surface S given by $x^2 + y^2 + z = 1$ for $z \geq 0$ is parametrized by $\Phi : D \rightarrow \mathbb{R}^3$, where D is the unit disk $u^2 + v^2 \leq 1$ and $\Phi(u, v) = (u, v, 1 - u^2 - v^2)$. S is also parametrized by $\Psi : R \rightarrow \mathbb{R}^3$, where R is the rectangle $[0, 1] \times [0, 2\pi]$ and $\Psi(r, \theta) = (r \cos(\theta), r \sin(\theta), 1 - r^2)$.

- A.** Φ is a one-to-one mapping of D onto $S = \Phi(D)$.
- B.** Ψ is a one-to-one mapping of R onto $S = \Psi(R)$.
- C.** $\Psi = \Phi \circ T$, where $T : R \rightarrow D$ is the polar coordinate transformation.
- D.** **A, B and C**
- *E.** **A and C**