

DUALITY

Ten Exercises for Bernd Sturmfels' Lectures
LMIPO, UC San Diego, March 22-23, 2010

1. Let P be the convex hull of three pairwise touching circles in \mathbb{R}^2 . Draw the dual convex body P^* . Now repeat the same with four spheres in \mathbb{R}^3 . Finally, what do you get when you take four pairwise touching circles in \mathbb{R}^3 ?
2. The curve $X = \{(x, y) \in \mathbb{R}^2 : x^4 + y^4 = 1\}$ is known as the *TV screen*. Determine the irreducible polynomial whose zero set is the dual curve X^* . Explain the results of our computation in terms of two dual norms on \mathbb{R}^2 .
3. Let X be the variety consisting of all $2 \times 2 \times 2$ -tensors of rank one. As a projective variety, X is the Segre embedding of $\mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$ into \mathbb{P}^7 . Compute the dual variety X^* . Verify the biduality $(X^*)^* = X$ for this example.
4. Draw a picture of the three-dimensional spectrahedron

$$P = \left\{ (x, y, z) \in \mathbb{R}^3 : \begin{pmatrix} 1 & x & 0 & x \\ x & 1 & y & 0 \\ 0 & y & 1 & z \\ x & 0 & z & 1 \end{pmatrix} \succeq 0 \right\}.$$

Also, compute the dual convex body P^* . Determine all faces of P and P^* .

5. Examine the trigonometric space curve which has the parametrization $x = \cos(\theta)$, $y = \cos(2\theta)$, $z = \sin(3\theta)$. Compute the convex hull of this curve.

6. Consider the problem of maximizing a linear function $ax + by + cz$ over the spectrahedron in Problem 4. Write down the dual semidefinite program and the critical equations. Derive an explicit formula, in terms of radicals in a , b and c , for the optimal value. Make sure your case distinction is complete.

7. Consider the problem of minimizing the trace of a positive semidefinite 10×10 -matrix whose off-diagonal entries are fixed to be random large integers. The optimal matrix has entries in a finite field extension of \mathbb{Q} . What ranks are possible for that matrix? What about the degree of the extension?

8. The semidefinite completion problem for a 4-cycle leads to the formula

$$\exists(x, y) \in \mathbb{R}^2 : \begin{pmatrix} u_{11} & u_{12} & x & u_{14} \\ u_{12} & u_{22} & u_{23} & y \\ x & u_{23} & u_{33} & u_{34} \\ u_{14} & y & u_{34} & u_{44} \end{pmatrix} \succeq 0.$$

This formula defines a convex cone $\mathcal{C}_{\mathcal{L}}$ in \mathbb{R}^8 . Compute the unique polynomial in the eight unknowns u_{ij} whose zero set is the algebraic boundary of $\mathcal{C}_{\mathcal{L}}$.

9. The polynomial $p(x) = 1 + x + x^2 + x^3 + x^4 + x^5 + x^6$ is non-negative on the real line. Find all possible representations of $p(x)$ as a sum of two squares. Does one of these representations involve only rational numbers? Show that the set of all SOS representations of $p(x)$ is a three-dimensional spectrahedron. Draw this spectrahedron and compute its analytic center.

10. For which values of the parameters a and b is the following polynomial non-negative on \mathbb{R}^2 ? In those cases, it is a sum of squares of polynomials.

$$f_{a,b}(x, y) = x^4 + y^4 + a(x^3 + y^2) + b(y^3 + x^2) + (a + b).$$

Draw that convex region \mathcal{C} in the (a, b) -plane. The *fiber* over $(a, b) \in \mathcal{C}$ is the spectrahedron whose points are the SOS representations of $f_{a,b}$. What are the various dimensions of the fibers as (a, b) ranges over the convex set \mathcal{C} ?