

## Supplement on the Monotone Class Theorem

In dealing with integrals, the following “functional” form of the Monotone Class Theorem is often useful.

**(1) Theorem.** Let  $\mathcal{K}$  be a collection of bounded real-valued functions on  $\Omega$  that is closed under the formation of products (i.e., if  $f, g \in \mathcal{K}$  then  $fg \in \mathcal{K}$ ), and let  $\mathcal{B}$  be the  $\sigma$ -algebra generated by  $\mathcal{K}$ . Let  $\mathcal{H} \supset \mathcal{K}$  be a vector space (over  $\mathbf{R}$ ) of bounded real-valued functions on  $\Omega$  such that (a)  $\mathcal{H}$  contains the constant functions and (b) if  $(f_n) \subset \mathcal{H}$  with  $\sup_n \sup_\omega |f_n(\omega)| < +\infty$  and if  $0 \leq f_1 \leq f_2 \leq \dots \leq f_n \leq \dots$ , then  $f := \lim_n f_n \in \mathcal{H}$ . Under these conditions,  $\mathcal{H}$  contains every bounded  $\mathcal{B}$ -measurable real-valued function on  $\Omega$ .

The proof of this theorem relies on the following

**(2) Lemma.** If  $\mathcal{H}$  is as in the statement of Theorem (1), then  $\mathcal{H}$  is closed under uniform convergence.

*Proof.* Suppose  $(f_n) \subset \mathcal{H}$  and  $f_n \rightarrow f$  uniformly on  $\Omega$ . (That is,  $\lim_n \sup_\omega |f_n(\omega) - f(\omega)| = 0$ .) By passing to a subsequence if necessary, we can arrange that  $\|f_{n+1} - f_n\|_\infty \leq 2^{-n}$ . Define  $g_n := f_n - 2^{1-n} + 2$ . Then  $g_n \in \mathcal{H}$  since  $\mathcal{H}$  is a vector space containing the constant functions, and

$$\sup_\omega |g_n(\omega)| \leq \sup_\omega |f_n(\omega)| + 2,$$

so the sequence  $(g_n)$  is uniformly bounded. Also,  $g_{n+1} - g_n = f_{n+1} - f_n + 2^{-n} \geq 0$  ( $n = 1, 2, \dots$ ) and  $g_1 = f_1 + 2^{-1} \geq 0$ . It follows that  $(g_n)$  is a (uniformly bounded) increasing sequence, with  $\lim_n g_n = \lim_n f_n + 2 = f + 2$ . But  $\lim_n g_n \in \mathcal{H}$ , hence so is  $f = \lim_n g_n - 2$ .  $\square$

*Proof of (1).* Owing to the closure properties of  $\mathcal{H}$  and the fact that every  $\mathcal{B}$ -measurable function is the pointwise limit of an increasing sequence of simple functions, it suffices to show that  $\mathcal{H}$  contains the indicator  $1_D$  of every  $D \in \mathcal{B}$ . Define  $\mathcal{L} := \{D \in \mathcal{B} : 1_D \in \mathcal{H}\}$ . It is easy to see that  $\mathcal{L}$  is a  $\lambda$ -system. We are going to show that  $\mathcal{L}$  contains a  $\pi$ -system  $\mathcal{P}$  generating  $\mathcal{B}$ . In view of the Monotone Class Theorem, this will imply  $\mathcal{L} \supset \mathcal{B}$ , whence  $\mathcal{L} = \mathcal{B}$ .

Let  $\mathcal{A}_0$  denote the algebra\* generated by  $\mathcal{K}$ ; since  $\mathcal{K}$  is already closed under products,  $\mathcal{A}_0$  is simply the linear span of  $\mathcal{K}$ . Consequently,  $\mathcal{A}_0 \subset \mathcal{H}$ . By Lemma (2), the uniform closure  $\mathcal{A}$  of  $\mathcal{A}_0$  is also contained in  $\mathcal{H}$ . Referring to the standard proof of Weierstrass' Theorem, we see that if  $f \in \mathcal{A}$ , then  $|f| \in \mathcal{A}$  as well. Consequently, if  $f, g \in \mathcal{A}$ , then  $f \vee g = [|f - g| + f + g]/2$  and  $f \wedge g = [f + g - |f - g|]/2$  are elements of  $\mathcal{A}$ . Now fix  $f \in \mathcal{A}$  and  $b \in \mathbf{R}$ . Then for each  $n \in \mathbf{N}$  the function  $\varphi_n := [n(f - b)^+] \wedge 1$  is an element of  $\mathcal{A}$ , hence an element of  $\mathcal{H}$ . As  $n \rightarrow \infty$ ,  $\varphi_n$  increases pointwise to  $1_{\{f > b\}}$ . Thus, since  $\mathcal{H}$  is closed under bounded monotone convergence,  $1_{\{f > b\}} \in \mathcal{H}$ ; this means that  $\{f > b\} \in \mathcal{L}$ . More generally, if  $\{f_1, f_2, \dots, f_m\}$  is a finite sequence of functions from  $\mathcal{A}$  and if  $\{b_1, b_2, \dots, b_m\}$  is a finite sequence of real numbers, then the function

$$g_n := \prod_{k=1}^m [n(f_k - b_k)^+] \wedge 1$$

---

\* An algebra (of functions) is a vector space that is also closed under the formation of products.

is an element of  $\mathcal{A}$ , and the sequence  $\{g_n\}$  increases boundedly and pointwise to  $\prod_{k=1}^m 1_{\{f_k > b_k\}}$ , which is therefore an element of  $\mathcal{H}$ , as before. This function is the indicator of the set  $B := \bigcap_{k=1}^m \{\omega : f_k(\omega) > b_k\}$ , so  $B \in \mathcal{L}$ . It follows that  $\mathcal{L}$  contains the  $\pi$ -system  $\mathcal{P}$  consisting of finite intersections of sets of the form  $f^{-1}(I)$ , where  $f \in \mathcal{A}$  and  $I \subset \mathbf{R}$  is an open right-halfline. Since  $\mathcal{K} \subset \mathcal{A}$ , the  $\sigma$ -algebra generated by  $\mathcal{P}$  is  $\mathcal{B}$  ( $= \sigma(\mathcal{K})$ ). We have constructed a  $\pi$ -system generating  $\mathcal{B}$  and contained in  $\mathcal{L}$ , as desired.  $\square$

**(3) Exercise.** Let  $(\Omega, \mathcal{B})$  be a measurable space. Let  $\mathcal{K}$  be a collection of bounded  $\mathcal{B}$ -measurable real-valued functions on  $\Omega$  such that  $\mathcal{B}$  is the  $\sigma$ -algebra generated by  $\mathcal{K}$ . Assume that  $\mathcal{K}$  is closed under the formation of products. Let  $P$  and  $Q$  be two probability measures on  $(\Omega, \mathcal{B})$  such that  $\int X dP = \int X dQ$  for all  $X \in \mathcal{K}$ . Prove that  $\int X dP = \int X dQ$  for every bounded  $\mathcal{B}$ -measurable random variable  $X$ . In particular,  $P = Q$  on  $\mathcal{B}$ . [Hint: Take  $\mathcal{H}$  to be the class of bounded  $\mathcal{B}$ -measurable random variables  $X$  such that  $\int X dP = \int X dQ$ , and apply Theorem (1).]

**(4) Example.** Let  $(X, \mathcal{M})$  and  $(Y, \mathcal{N})$  be measurable spaces, and recall that  $\mathcal{M} \otimes \mathcal{N}$  denotes the product  $\sigma$ -algebra on the cartesian product  $X \times Y$ . More precisely,  $\mathcal{M} \otimes \mathcal{N}$  is the  $\sigma$ -algebra on  $X \times Y$  generated by the projections  $\pi_1, \pi_2$ , where  $\pi_1(x, y) = x$  and  $\pi_2(x, y) = y$  for  $(x, y) \in X \times Y$ . Let  $\mathcal{K}$  denote the set of functions of the form  $(x, y) \mapsto f(x)g(y)$ , where  $f$  (resp.  $g$ ) is a bounded real-valued  $\mathcal{M}$ -measurable (resp.  $\mathcal{N}$ -measurable) function on  $X$  (resp.  $Y$ ). Clearly the  $\sigma$ -algebra generated by  $\mathcal{K}$  is just  $\mathcal{M} \otimes \mathcal{N}$ . Now let  $\mathcal{H}$  be the set of bounded real-valued  $\mathcal{M} \otimes \mathcal{N}$ -measurable functions  $h$  such that  $x \mapsto h(x, y)$  is  $\mathcal{M}$ -measurable for each fixed  $y \in Y$ . It is a simple matter to check that  $\mathcal{H}$  is a vector space satisfying the conditions of Theorem (1). As a consequence of that result,  $\mathcal{H}$  is precisely the class of *all* bounded real-valued  $\mathcal{M} \otimes \mathcal{N}$ -measurable functions. This is an alternative proof of one of the measurability assertions in Tonelli's theorem, at least for bounded functions. A truncation argument reduces the general case to the bounded case.

**(5) Exercise.** Let  $(\Omega, \mathcal{B}, P)$  be a probability space, and let  $X$  and  $Y$  be random variables defined on  $(\Omega, \mathcal{B})$ . Suppose that

$$(6) \quad E[f(X)g(Y)] = E[f(X)g(X)]$$

for every pair  $(f, g)$  of bounded continuous functions from  $\mathbf{R}$  to  $\mathbf{R}$ . Prove that  $P[X = Y] = 1$ . [Hint: Use Theorem (1) to show that  $E[h(X, Y)] = E[h(X, X)]$  for every bounded  $\mathcal{B}(\mathbf{R}^2)$ -measurable real-valued function  $h$ . Then consider  $h = 1_\Delta$ , where  $\Delta = \{(x, x) : x \in \mathbf{R}\}$  is the "diagonal" in  $\mathbf{R}^2$ .]

The following variant of Theorem (1) is sometimes useful. The proof is quite similar to that of Theorem (1), and so is omitted.

**(7) Theorem.** *Let  $\mathcal{C}$  be an algebra of bounded real-valued functions on  $\Omega$  that contains the constant functions, and let  $\mathcal{B}$  be the  $\sigma$ -algebra generated by  $\mathcal{C}$ . Let  $\mathcal{H} \supset \mathcal{C}$  be a set of bounded real-valued functions on  $X$  that is closed under bounded monotone convergence and uniform convergence. Under these conditions,  $\mathcal{H}$  contains every bounded  $\mathcal{B}$ -measurable real-valued function on  $\Omega$ .*

As an application of Theorem (7), you are asked to prove a functional form of Exercise 5 from Chapter 2.

**(8) Exercise.** Let  $(S, d)$  be a metric space, let  $BC(S)$  denote the class of bounded continuous real-valued functions on  $S$ , and let  $\mathcal{B}(S)$  denote the Borel  $\sigma$ -field on  $S$ ; thus  $\mathcal{B}(S)$  is the  $\sigma$ -field generated by the open subsets of  $S$ .

- (a) Verify that  $\mathcal{B}(S)$  is the  $\sigma$ -algebra generated by  $BC(S)$ .
- (b) Let  $\mu$  be a probability measure on  $(S, \mathcal{B}(S))$  and let  $f$  be a bounded real-valued  $\mathcal{B}(S)$ -measurable function. Using Theorem (7), show that for each  $\epsilon > 0$  there exists  $g \in BC(S)$  such that  $\int |f - g| d\mu \leq \epsilon$ . [Hint: Take  $\mathcal{H}$  to be the class of bounded real-valued  $\mathcal{B}(S)$ -measurable functions for which the asserted approximation property holds, and take  $\mathcal{C} = BC(S)$ . Show that  $\mathcal{H}$  has the required closure properties for Theorem (7) to apply.]

**(9) Example.** Let  $\{X_t : t \in \mathbf{T}\}$  be a collection of random variables, indexed by  $\mathbf{T}$ , defined on a common measurable space  $(\Omega, \mathcal{B})$ . Let  $\mathcal{X} \subset \mathcal{B}$  denote the  $\sigma$ -field generated by  $\{X_t : t \in \mathbf{T}\}$ .

**Claim:** If  $F : \Omega \rightarrow \mathbf{R}$  is  $\mathcal{X}$ -measurable, then there is a sequence  $\{t_1, t_2, \dots\} \subset \mathbf{T}$  and a  $\mathcal{B}(\mathbf{R}^{\mathbf{N}})$ -measurable function  $f : \mathbf{R}^{\mathbf{N}} \rightarrow \mathbf{R}$  such that

$$(10) \quad F(\omega) = f(X_{t_1}(\omega), X_{t_2}(\omega), \dots), \quad \forall \omega \in \Omega.$$

To see the Claim apply Theorem (1) with  $\mathcal{K}$  equal to the class of functions of the form

$$\omega \mapsto \prod_{k=1}^n f_k(X_{t_k}(\omega)),$$

where  $n \in \mathbf{N}$ ,  $t_k \in \mathbf{T}$ , and each  $f_k : \mathbf{R} \rightarrow \mathbf{R}$  is bounded and  $\mathcal{B}(\mathbf{R})$ -measurable. Observe that  $\sigma(\mathcal{K}) = \mathcal{X}$ . Take  $\mathcal{H}$  to be the class of bounded  $\mathcal{X}$ -measurable functions for which a representation like (10) holds. It should be clear that  $\mathcal{H}$  is a vector space containing the constant functions. Let  $\{F_n\}_{n \in \mathbf{N}}$  be a uniformly bounded increasing sequence of elements of  $\mathcal{H}$ . Since a countable union of countable sets is itself countable, we may suppose that there is a single sequence  $\{t_1, t_2, \dots\}$  such that

$$F_n(\omega) = f_n(X_{t_1}(\omega), X_{t_2}(\omega), \dots), \quad \forall n \in \mathbf{N}, \omega \in \Omega,$$

where each  $f_n$  is bounded and  $\mathcal{B}(\mathbf{R}^{\mathbf{N}})$ -measurable. Since  $\{F_n\}$  is uniformly bounded, we can suppose that there is a constant  $M$  such that  $|f_n(\mathbf{x})| \leq M$  for all  $n \in \mathbf{N}$  and all  $\mathbf{x} \in \mathbf{R}^{\mathbf{N}}$ . Define  $f : \mathbf{R}^{\mathbf{N}} \rightarrow \mathbf{R}$  by

$$f(\mathbf{x}) := \liminf_{n \rightarrow \infty} f_n(\mathbf{x}), \quad \mathbf{x} \in \mathbf{R}^{\mathbf{N}}.$$

Then  $f$  is bounded in magnitude by  $M$  and is  $\mathcal{B}(\mathbf{R}^{\mathbf{N}})$ -measurable. Moreover, because  $F_n(\omega)$  increases with  $n$ ,

$$F(\omega) := \lim_n F_n(\omega) = \lim_n \inf_n F_n(\omega) = \lim_n \inf_n f_n(X_{t_1}(\omega), X_{t_2}(\omega), \dots) = f(X_{t_1}(\omega), X_{t_2}(\omega), \dots),$$

so  $F$  admits a representation as in (10). That is,  $\mathcal{H}$  is closed under bounded monotone convergence. By Theorem (1),  $\mathcal{H}$  contains every bounded  $\mathcal{X}$ -measurable real-valued function on  $\Omega$ . This proves the Claim.