FIFTY-SIXTH ANNUAL
WILLIAM LOWELL PUTNAM MATHEMATICAL COMPETITION

Saturday, December 2, 1995 Examination A

A-1. Let S be a set of real numbers which is closed under multiplication (that is, if a and b are in S, then so is ab). Let T and U be disjoint subsets of S whose union is S. Given that the product of any *three* (not necessarily distinct) elements of T is in T and that the product of any three elements of U is in U, show that at least one of the two subsets T, U is closed under multiplication.

A-2. For what pairs (a, b) of positive real numbers does the improper integral

$$\int_b^\infty \left(\sqrt{\sqrt{x+a} - \sqrt{x}} - \sqrt{\sqrt{x} - \sqrt{x-b}} \right) dx$$

converge?

A-3. The number $d_1 d_2 \ldots d_9$ has nine (not necessarily distinct) decimal digits. The number $e_1 e_2 \ldots e_9$ is such that each of the nine 9-digit numbers formed by replacing just one of the digits d_i in $d_1 d_2 \ldots d_9$ by the corresponding digit e_i $(1 \leq i \leq 9)$ is divisible by 7. The number $f_1 f_2 \ldots f_9$ is related to $e_1 e_2 \ldots e_9$ is the same way: that is, each of the nine numbers formed by replacing one of the e_i by the corresponding f_i is divisible by 7. Show that, for each i, $d_i - f_i$ is divisible by 7. [For example, if $d_1 d_2 \ldots d_9 = 199501996$, then e_6 may be 2 or 9, since 199502996 and 199509996 are multiples of 7.]

A-4. Suppose we have a necklace of n beads. Each bead is labeled with an integer and the sum of all these labels is $n - 1$. Prove that we can cut the necklace to form a string whose consecutive labels $x_1, x_2, \ldots, x_n$ satisfy

$$\sum_{i=1}^{k} x_i \leq k - 1 \qquad \text{for } k = 1, 2, \ldots, n.$$

A-5. Let $x_1, x_2, \ldots, x_n$ be differentiable (real-valued) functions of a single variable t which satisfy

$$\begin{aligned}
\frac{dx_1}{dt} &= a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \\
\frac{dx_2}{dt} &= a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n \\
&\vdots \qquad\quad \vdots \qquad \vdots \\
\frac{dx_n}{dt} &= a_{n1}x_1 + a_{n2}x_2 + \cdots + a_{nn}x_n
\end{aligned}$$

for some constants $a_{ij} > 0$. Suppose that for all i, $x_i(t) \to 0$ as $t \to \infty$. Are the functions $x_1, x_2, \ldots, x_n$ necessarily linearly dependent?

A-6. Suppose that each of n people writes down the numbers 1, 2, 3 in random order in one column of a $3 \times n$ matrix, with all orders equally likely and with the orders for different columns independent of each other. Let the row sums a, b, c of the resulting matrix be rearranged (if necessary) so that $a \leq b \leq c$. Show that for some $n \geq 1995$, it is at least four times as likely that both $b = a + 1$ and $c = a + 2$ as that $a = b = c$.

FIFTY-SIXTH ANNUAL
WILLIAM LOWELL PUTNAM MATHEMATICAL COMPETITION

Saturday, December 2, 1995 Examination B

B-1. For a partition π of $\{1,2,3,4,5,6,7,8,9\}$, let $\pi(x)$ be the number of elements in the part containing x. Prove that for any two partitions π and π', there are two distinct numbers x and y in $\{1,2,3,4,5,6,7,8,9\}$ such that $\pi(x) = \pi(y)$ and $\pi'(x) = \pi'(y)$. [A *partition* of a set S is a collection of disjoint subsets (parts) whose union is S.]

B-2. An ellipse, whose semi-axes have lengths a and b, rolls without slipping on the curve $y = c\sin(x/a)$. How are a, b, c related, given that the ellipse completes one revolution when it traverses one period of the curve?

B-3. To each positive integer with n^2 decimal digits we associate the determinant of the matrix obtained by writing the digits in order across the rows. For example, for $n = 2$, to the integer 8617 we associate $\det\begin{pmatrix} 8 & 6 \\ 1 & 7 \end{pmatrix} = 50$. Find, as a function of n, the sum of all the determinants associated with n^2-digit integers. (Leading digits are assumed to be nonzero; for example, for $n = 2$, there are 9000 determinants.)

B-4. Evaluate

$$\sqrt[8]{2207 - \frac{1}{2207 - \frac{1}{2207 - \ldots}}}.$$

Express your answer in the form $(a + b\sqrt{c})/d$, where a, b, c, d are integers.

B-5. A game starts with four heaps of beans, containing 3,4,5 and 6 beans. The two players move alternately. A move consists of taking *either*

a. one bean from a heap, provided at least two beans are left behind in that heap, or

b. a complete heap of two or three beans.

The player who takes the last heap wins. To win the game, do you want to move first or second? Give a winning strategy.

B-6. For a positive real number α, define

$$S(\alpha) = \{\lfloor n\alpha \rfloor : n = 1, 2, 3, \ldots\}.$$

Prove that $\{1, 2, 3, \ldots\}$ cannot be expressed as the disjoint union of three sets $S(\alpha)$, $S(\beta)$ and $S(\gamma)$. [As usual, $\lfloor x \rfloor$ is the greatest integer $\leq x$.]