

The impact of bootstrap methods on time series analysis

Dimitris N. Politis*

1 Introduction: the sample mean of a time series

Let X_1, \dots, X_n be an observed stretch from a strictly stationary time series $\{X_t, t \in \mathbf{Z}\}$; the assumption of stationarity implies that the joint probability law of $(X_t, X_{t+1}, \dots, X_{t+k})$ does not depend on t for any $k \geq 0$. Assume also that the time series is weakly dependent, i.e., the collection of random variables $\{X_t, t \leq 0\}$ is approximately independent of $\{X_t, t \geq k\}$ when k is large enough. An example of a weak dependence structure is given by m -dependence under which $\{X_t, t \leq 0\}$ is (exactly) independent of $\{X_t, t \geq k\}$ whenever $k > m$; independence is just the special case of 0-dependence.

Due to the dependence between the observations, even the most basic calculations involved in applied statistical work suddenly become challenging; an elementary such example has to do with estimating the unknown mean $\mu = EX_t$ of the time series. The sample mean $\bar{X}_n = n^{-1} \sum_{t=1}^n X_t$ is the obvious estimator; however—and here immediately the difficulties crop up—it is not the most efficient. To see why, consider the regression $X_t = \mu + \epsilon_t$, that also serves as a definition for the ϵ_t process; it is apparent that \bar{X}_n is the Ordinary Least Squares Estimator of μ in this model. However, because of the dependence in the errors ϵ_t , the Best Linear Unbiased Estimator (BLUE) of μ is instead obtained by a Generalized Least Squares argument; consequently, $\hat{\mu}_{BLUE} = (\mathbf{1}'\Gamma_n^{-1}\mathbf{1})^{-1}\mathbf{1}'\Gamma_n^{-1}\mathbf{X}$, where $\mathbf{X} = (X_1, \dots, X_n)'$, $\mathbf{1} = (1, \dots, 1)'$, and Γ_n is the (unknown) covariance matrix of the vector \mathbf{X} with i, j element given by $\gamma(i - j) = Cov(X_i, X_j)$.

* Professor of Mathematics and Adjunct Professor of Economics, University of California, San Diego, La Jolla, CA 92093-0112, USA; e-mail: dpolitis@ucsd.edu. Research partially supported by NSF grant DMS-01-04059. Many thanks are due to Prof. E. Paparoditis (Cyprus) and J. Romano (Stanford) for constructive comments on this manuscript.

It is immediate that $\hat{\mu}_{BLUE}$ is a weighted average of the X_t data, i.e., $\hat{\mu}_{BLUE} = \sum_{i=1}^n w_i X_i$; the weights w_i are a function of the (unknown) covariances $\gamma(s)$, $s = 0, 1, \dots, n-1$. For example, under the simple AR(p), i.e., autoregressive of order p , model:

$$(X_t - \mu) = \phi_1(X_{t-1} - \mu) + \dots + \phi_p(X_{t-p} - \mu) + Z_t, \quad (1)$$

where $Z_t \sim$ i.i.d. $(0, \tau^2)$, it is easy to see that $w_j = w_{n-j+1}$ for $j = 1, \dots, p$, $w_{p+1} = w_{p+2} = \dots = w_{n-p}$, and $\sum_{i=1}^n w_i = 1$; calculating w_1, \dots, w_p in terms of $\gamma(\cdot)$ is feasible but cumbersome—see Fuller (1996). For instance, in the special case of $p = 1$, we can calculate that $w_1 = w_2/(1 - \rho(1))$ where $\rho(s) = \gamma(s)/\gamma(0)$ is the autocorrelation sequence.

Nevertheless, it is apparent that, at least in the AR(p) model above, the difference of the w_i weights from the constant weights in \bar{X}_n is due mostly to the end effects, i.e., the first and last p weights. Thus, it may be reasonable to conjecture that, for a large sample size n , those end effects will have a negligible contribution. This is in fact true; as shown in Grenander and Rosenblatt (1957) for a large class of weakly dependent processes (not limited to the AR(p) models) the sample mean \bar{X}_n is asymptotically efficient. Therefore, estimating μ by our familiar, easy-to-use, sample mean \bar{X}_n is fortuitously justified.

The problem, however, is further complicated if one wants to construct an estimate of the standard error of \bar{X}_n . To see this, let $\sigma_n^2 = Var(\sqrt{n}\bar{X}_n)$, and note that $n\sigma_n^2$ is just the sum of all the elements of the matrix Γ_n , i.e., $\sigma_n^2 = \sum_{s=-n}^n (1 - |s|/n)\gamma(s)$. Under usual regularity conditions,

$$\sigma_\infty^2 := \lim_{n \rightarrow \infty} \sigma_n^2 = \sum_{s=-\infty}^{\infty} \gamma(s) = 2\pi f(0),$$

where $f(w) = (2\pi)^{-1} \sum_{s=-\infty}^{\infty} e^{iws} \gamma(s)$, for $w \in [-\pi, \pi]$, is the spectral density function of the $\{X_t\}$ process. It is apparent now that the simple task of standard error estimation is highly nontrivial under dependence, as it amounts to nonparametric estimation of the spectral density at the origin.

One may think that estimating γ and plugging into the above formula would give a consistent estimator of the infinite sum $\sum_{s=-\infty}^{\infty} \gamma(s)$. Estimators of $\gamma(s)$ for $|s| < n$ can certainly be constructed; $\tilde{\gamma}(s) = (n - s)^{-1} \sum_{t=1}^{n-s} (X_t - \bar{X}_n)(X_{t+s} - \bar{X}_n)$ and $\hat{\gamma}(s) = n^{-1} \sum_{t=1}^{n-s} (X_t - \bar{X}_n)(X_{t+s} - \bar{X}_n)$ are the usual candidates. Note that $\tilde{\gamma}(s)$ is unbiased but becomes quite unreliable for large s because of increasing variability; at the extreme case of $s = n - 1$, there is no averaging at all in $\tilde{\gamma}(s)$. Thus, $\hat{\gamma}(s)$ is preferable for two reasons: (a) it trades in some bias for a huge variance reduction

by using the assumption of weak dependence (which implies that $\gamma(s) \rightarrow 0$ as $s \rightarrow \infty$) to shrink the $\tilde{\gamma}(s)$ towards zero for large s ; and (b) $\hat{\gamma}(s)$ is a nonnegative definite sequence, i.e., it guarantees that $\sum_{|s|<n} \hat{\gamma}(s) \geq 0$. To understand the last claim, note that $\sum_{|s|<n} \hat{\gamma}(s) = 2\pi T(0)$, where $T(w)$ is the periodogram defined by

$$T(w) = (2\pi n)^{-1} \left| \sum_{s=1}^n e^{iws} (X_s - \bar{X}_n) \right|^2$$

which, however, is well-known to be inconsistent as an estimator of $f(w)$; although the bias of $T(w)$ is negligible (of order $O(1/n)$), its variance is approximately constant and does not approach zero—see e.g. Brockwell and Davis (1991). Consequently, the plug-in idea $\sum_{|s|<n} \hat{\gamma}(s)$ is inconsistent as well; intuitively, it is seen that $\hat{\gamma}(s)$ is still unreliable for large s and more shrinking towards zero is necessary.

The above observations suggest two different ways of achieving consistent spectral estimation:

- A. Split the data series X_1, \dots, X_n into (overlapping) blocks of size b , i.e., (X_1, \dots, X_b) , (X_2, \dots, X_{b+1}) , etc. Calculate the periodogram $T_i(w)$ from block (X_i, \dots, X_{i+b-1}) , for $i = 1, \dots, q$ (where $q = n - b + 1$), and then let $\bar{T}(w) = q^{-1} \sum_{i=1}^q T_i(w)$ be the average of those short periodograms. It follows that the bias of $\bar{T}(w)$ is equal to the bias of each of the $T_i(w)$, i.e., it is of order $O(1/b)$, while its variance¹ is $O(b/n)$ due to the averaging; thus, consistency of $\bar{T}(w)$ ensues provided

$$b \rightarrow \infty \text{ as } n \rightarrow \infty \text{ but with } b/n \rightarrow 0. \quad (2)$$

This blocking idea was actually one of the very first spectral estimation schemes proposed in a pioneering paper by Bartlett (1946).

- B. Since more shrinking of $\hat{\gamma}(s)$ towards zero is advisable for the purposes of convergence of their sum, it comes as no surprise that a popular method of estimating $f(w)$ is given by means of truncated/weighted sums:

$$\hat{f}_{b,\lambda}(w) = (2\pi)^{-1} \sum_{s=-b}^b \lambda(s/b) e^{iws} \hat{\gamma}(s), \quad (3)$$

¹To see that $\text{Var}(\bar{T}(w)) = O(b/n)$, note that periodograms calculated from nonoverlapping blocks are approximately independent; there are approximately n/b such nonoverlapping blocks within the data stretch X_1, \dots, X_n . By a Cauchy-Schwarz argument, it is immediate that the variance of $\bar{T}(w)$ can not be bigger than the variance of the average of just those n/b approximately independent, nonoverlapping periodograms which is $O(b/n)$.

where the lag-window λ is some (well-chosen) function on $[-1, 1]$. Consistency is achieved again under (2); in particular, the condition $b = o(n)$ effectively keeps the unreliable $\hat{\gamma}(s)$ —the ones for large s —from entering in and spoiling the convergence of the sum in eq. (3).

Interestingly, the above two methods of spectral estimation turn out to be equivalent to each other—as well as to the third popular method that entails smoothing the periodogram; see Brockwell and Davis (1991). For example, Bartlett’s $\bar{T}(w)$ is approximately equivalent to $\hat{f}_{b,\lambda}(w)$ using the triangular lag-window $\lambda_B(x) = 1 - |x|$. Although the constant lag-window $\lambda_C(x) = 1$ also leads to consistent estimation under (2), improved accuracy is achieved via the family of flat-top lag-windows defined in Politis and Romano (1995) the simplest representative of which is the trapezoidal lag-window $\lambda_T(x) = \min\{1, 2(1 - |x|)\}$ that can be thought of as a compromise between λ_C and λ_B .

2 Block resampling and subsampling

As is well known, the bootstrap scheme pioneered in the seminal paper of Efron (1979) was geared towards independent data. As a matter of fact, it was quickly recognized by Singh (1981) that if one applies the i.i.d. bootstrap to data that are dependent, inconsistency follows. Because of the total data-‘scrambling’ induced by the i.i.d. bootstrap all dependence information is lost. Consequently, the i.i.d. bootstrap estimate of σ_n^2 typically does not converge to the correct limit given by $\sigma_\infty^2 = \sum_{s=-\infty}^{\infty} \gamma(s)$; rather, it converges to $\gamma(0)$, i.e., the i.i.d. bootstrap fails to give valid standard error estimates in the dependent case.

2.1 Subsampling

Subsampling is probably one of the most intuitive methods of valid statistical inference. To see why subsampling works in the time series case, consider the aforementioned consistent estimator of σ_∞^2 due to Bartlett, i.e., $2\pi\bar{T}(0)$ or equivalently $2\pi\hat{f}_{b,\lambda_B}(0)$. Recall that $2\pi\hat{f}_{b,\lambda_B}(0) = \sum_{s=-b}^b (1 - |s|/b)\hat{\gamma}(s)$ which is actually a simple, plug-in estimator of $\sigma_b^2 = \text{Var}(\sqrt{b}\sum_{i=1}^b X_i) = \sum_{s=-b}^b (1 - |s|/b)\gamma(s)$. Intuitively, $2\pi\hat{f}_{b,\lambda_B}(0)$ is estimating σ_b^2 and not σ_n^2 ; consistency of $2\pi\hat{f}_{b,\lambda_B}(0)$ is achieved only because both σ_b^2 and σ_n^2 tend to σ_∞^2 under (2)—and therefore also $\sigma_b^2 - \sigma_n^2 \rightarrow 0$.

In other words, it seems that we cannot directly estimate σ_n^2 or σ_∞^2 , and we have to contend ourselves with estimating σ_b^2 . But an immediate,

empirical estimator of $\sigma_b^2 = \text{Var}(\sqrt{b} \sum_{i=1}^b X_i)$ can be constructed from the many size- b sample means that can be extracted from our data X_1, \dots, X_n . To formalize this notion, let $\bar{X}_{i,b} = \sum_{t=i}^{i+b-1} X_t$ be the sample mean of the i th block, and let $\hat{\sigma}_{b,SUB}^2$ denote the sample variance of the (normalized) subsample values $\sqrt{b}\bar{X}_{i,b}$ for $i = 1, \dots, q$. The simple estimator $\hat{\sigma}_{b,SUB}^2$ is the subsampling estimator of variance; it is consistent for σ_∞^2 under (2) which is not surprising in view of that fact that it, too, is equivalent to the ubiquitous Bartlett estimator!

Nevertheless, the beauty of the subsampling methodology is its extreme generality: let $\hat{\theta}_n = \hat{\theta}_n(X_1, \dots, X_n)$ be an arbitrary statistic that is consistent for a general parameter θ at rate a_n , i.e., for large n , $a_n(\hat{\theta}_n - \theta)$ tends to some well-defined asymptotic distribution J ; the rate a_n does not have to equal \sqrt{n} , and the distribution J does not have to be normal—we do not even need to know its shape, just that it exists. Let $\hat{\theta}_{i,b} = \hat{\theta}_b(X_i, \dots, X_{i+b-1})$ be the subsample value of the statistic computed from the i th block. The subsampling estimator of J is $\hat{J}_{b,SUB}$ defined as the empirical distribution of the normalized (and centered) subsample values $a_b(\hat{\theta}_{i,b} - \hat{\theta}_n)$ for $i = 1, \dots, q$.

The consistency of $\hat{J}_{b,SUB}$ for general statistics under minimal conditions—basically involving a weak dependence condition and (2)—was shown in Politis and Romano (1992c, 1994b); consequently, confidence intervals for θ can immediately be formed using the quantiles of $\hat{J}_{b,SUB}$ instead of the quantiles of the (unknown) J . A closely related early work is the paper by Sherman and Carlstein (1996) where the subsampling distribution of the (unnormalized) subsample values is put forth as a helpful diagnostic tool.

If a variance estimator for $a_n\hat{\theta}_n$ is sought, it can be constructed by the sample variance of the normalized subsample values $a_b\hat{\theta}_{i,b}$ for $i = 1, \dots, q$; consistency of the subsampling estimator of variance for general statistics was shown by Carlstein (1986) under some uniform integrability conditions. It should be pointed out that if there is no dependence present, then the ordering of the data is immaterial; the data can be permuted with no information loss. Thus, in the i.i.d. case there are $\binom{n}{b}$ subsamples of size b from which subsample statistics can be recomputed in which case subsampling is equivalent to the well-known delete- d jackknife (with $d = n - b$) of Wu (1986) and Shao and Wu (1989); see Politis, Romano and Wolf (1999) for more details and extensive list of references.

Subsampling in the i.i.d. case is also intimately related to the i.i.d. bootstrap with smaller resample size b , i.e., where our statistic is re-computed from X_1^*, \dots, X_b^* drawn i.i.d. from the empirical distribution of the data X_1, \dots, X_n ; the only difference is that subsampling draws b values *without*

replacement from the dataset X_1, \dots, X_n while the bootstrap draws *with* replacement.

The possibility of drawing an arbitrary resample size was noted as early on as in Bickel and Freedman (1981); nevertheless, the research community focused for some time on the case of resample size n since that choice leads to improved performance, i.e., higher-order accuracy, in the sample mean case—cf. Singh (1981). Nevertheless, in instances where the i.i.d. bootstrap fails, it was observed by many researchers, most notably K. Athreya, J. Bretagnolle, M. Arcones and E. Giné, that bootstrap consistency can be restored by taking a smaller resample size b satisfying (2); see Politis et al. (1999) for references but also note the early paper by Swanepoel (1986).

As a matter of fact, the i.i.d. bootstrap with resample size b immediately inherits the general validity of subsampling in the case where $b = o(\sqrt{n})$ in which case the question with/without replacement is immaterial; see Politis and Romano (1992c, 1993) or Politis et al. (1999, Corollary 2.3.1). However, to relax the condition $b = o(\sqrt{n})$ to $b = o(n)$ some extra structure is required that has to be checked on a case-by-case basis; see Bickel et al. (1997).

Finally, note that in the case of the sample mean of i.i.d. data (with finite variance), the bootstrap with resample size n outperforms both subsampling and/or the bootstrap with smaller resample size, as well as the asymptotic normal approximation. Although the performance of subsampling can be boosted by the use of extrapolation and interpolation techniques—see e.g., Booth and Hall (1993), Politis and Romano (1995), Bickel et al. (1997), Bertail and Politis (2001)—the understanding has been that subsampling sacrifices some accuracy for its extremely general applicability; this was also the viewpoint adopted in Politis et al. (1999, ch. 10). Nevertheless, some very recent results of Sakov and Bickel (2000) and Arcones (2001) indicate that this is not always the case: for the sample median, the i.i.d. bootstrap with resample size $b = o(n)$ is the one that leads to improved accuracy and *not* the bootstrap with resample size n ; an analogous result is expected to hold for subsampling.

2.2 Block resampling

In the last section we saw the connection of subsampling in the i.i.d. case to the delete- d jackknife. Recall that the standard delete-1 jackknife is commonly attributed to Tukey (1958) and to Quenouille (1949, 1956). Interestingly, Quenouille’s work was focused on blocking methods in a time series context, and thus it was the precursor of (block) subsampling; of course, blocking methods for estimation in time series go back to Bartlett (1946) as

previously mentioned.

After Carlstein’s (1986) subseries variance estimator—as well as Hall’s (1985) re-tiling ideas for spatial data—the time was ripe for Künsch’s (1989) introduction of the block bootstrap where, instead of re-computing the statistic $\hat{\theta}_b$ on the smaller blocks of the type $B_i = (X_i, \dots, X_{i+b-1})$, a new bootstrap pseudo-series X_1^*, \dots, X_l^* is created (with $l = kb \simeq n$) by joining together $k (= \lceil n/b \rceil)$ blocks chosen randomly (and with replacement) from the set $\{B_1, \dots, B_{n-b+1}\}$; then the statistic can be re-computed from the new, full-size pseudo-series X_1^*, \dots, X_l^* .

Thus, ten years after Efron’s (1979) pioneering paper, Künsch’s (1989) breakthrough indicated the starting point for a focus of research activity on this new area of block resampling. The closely related work by Liu and Singh (1992) which was available as a preprint in the late 80s was also quite influential at that point because of its transparent and easily generalizable approach.

Examples of some early work on block resampling for stationary time series include the blocks-of-blocks bootstrap of Politis and Romano (1992a) that extended the applicability of the block bootstrap to parameters associated with the whole infinite-dimensional probability law of the series, the circular bootstrap of Politis and Romano (1992b) in which the bootstrap distribution is automatically centered correctly, and the stationary bootstrap of Politis and Romano (1994a) that joins together blocks of random length—having a geometric distribution with mean b —and thus generates bootstrap sample paths that are stationary series themselves.

As in the i.i.d. case, the validity of block bootstrap methods must be checked on a case-by-case basis, and it is usually facilitated by an underlying asymptotic normality; typically, a condition such as (2) is required. For instance, the block bootstrap estimator of the distribution of $\sqrt{n}(\bar{X}_n - \mu)$ is essentially consistent only if the latter tends to a Gaussian limit, a result that parallels the i.i.d. situation; see Giné and Zinn (1990) and Radulovic (1996). To see why, note that in the sample mean case the block bootstrap distribution is really a (rescaled) k -fold convolution of the subsampling distribution $\hat{J}_{b,SUB}$ with itself; since $k = \lceil n/b \rceil \rightarrow \infty$ under (2), it is intuitive that the block bootstrap distribution has a strong tendency towards Gaussianity.

Interestingly, the block bootstrap estimator $\hat{\sigma}_{b,BB}^2$ of $Var(\sqrt{n}\bar{X}_n)$ is identical to the subsampling estimator $\hat{\sigma}_{b,SUB}^2$ which in turn is equivalent to the Bartlett estimator $2\pi\hat{f}_{b,\lambda_B}(0)$; recall that the latter is suboptimal due to its rather large bias of order $O(1/b)$. In addition, the circular and stationary bootstrap estimators of $Var(\sqrt{n}\bar{X}_n)$ suffer from a bias of the same order

of magnitude; see Lahiri (1999). However, if variance estimation is our objective, we can construct a highly accurate estimator via a linear combination of block resampling/subsampling estimators, in the same way that the trapezoidal lag-window spectral estimator \hat{f}_{b,λ_T} can be gotten by a linear combination of two Bartlett estimators. The simplest such proposal is to define

$$\tilde{\sigma}_b^2 = 2\hat{\sigma}_{b,BB}^2 - \hat{\sigma}_{b/2,BB}^2$$

where b is now an even integer. The estimator $\tilde{\sigma}_b^2$ has bias (and MSE) that is typically orders of magnitude less than those of $\hat{\sigma}_{b,BB}^2$; as a matter of fact, $\tilde{\sigma}_b^2 \simeq 2\pi\hat{f}_{b,\lambda_T}(0)$ which, as previously alluded to, enjoys some optimality properties. For example, if the covariance $\gamma(s)$ has an exponential decay—as is the case in stationary ARMA models—, then the bias of $\tilde{\sigma}_b^2$ becomes $O(e^{-b})$; see Politis and Romano (1995), Politis et al. (1999, Ch. 10.5) or Bertail and Politis (2001).

A theoretical disadvantage of $\tilde{\sigma}_b^2$ is that it is not almost surely non-negative—although its fast convergence to a nonnegative limit practically alleviates this problem; this comes as no surprise since the trapezoidal lag-window corresponds to a higher-order (actually, *infinite-order*) smoothing kernel. Nevertheless, the implication is that $\tilde{\sigma}_b^2$ is not associated with a bootstrap distribution for \bar{X}_n , as is the case with $\hat{\sigma}_{b,BB}^2$.

In order to come up with a method that gives improved variance estimation while at the same time being able to produce bootstrap pseudo-series and a bootstrap distribution for \bar{X}_n the following key idea may be employed: the B_i blocks may be *tapered*, i.e., their end-points are shrunk towards a target value, before being concatenated to form a bootstrap pseudo-series; this is the tapered block bootstrap method of Paparoditis and Politis (2001a, 2002b) that indeed leads to a variance estimator $\hat{\sigma}_{b,TBB}^2$ that is more accurate than $\hat{\sigma}_{b,BB}^2$. In particular, the bias of $\hat{\sigma}_{b,TBB}^2$ is of order $O(1/b^2)$. Although $\hat{\sigma}_{b,BB}^2$ is typically less accurate than $\tilde{\sigma}_b^2$, the tapered block bootstrap constructs an estimator of the whole distribution for \bar{X}_n (not just its variance), and this estimator is more accurate than its (untapered) block bootstrap analog. Intuitively, this preferential treatment (tapering) of the block edges is analogous to the weighting of the boundary points in constructing $\hat{\mu}_{BLUE}$ as discussed in the Introduction.

2.3 Block size choice and some further issues

Recall that the block size b must be chosen in order to practically implement any of the block resampling/subsampling methods. By the equivalence of

$\hat{\sigma}_{b,BB}^2$ and $\hat{\sigma}_{b,SUB}^2$ to the Bartlett spectral estimator $2\pi\hat{f}_{b,\lambda_B}(0)$ it is apparent that choosing b is tantamount to the difficult problem of bandwidth choice in nonparametric smoothing problems. As in the latter problem, here too there are two main approaches:

- (a) *Cross-validation.* A promising subsampling/cross-validation approach for block size choice proposed by Hall et al. (1995).
- (b) *Plug-in methods.* This approach entails working out an expression for the optimal (with respect to some criterion) value of b , and then estimating/plugging-in all unknown parameters in that expression.

In order for the plug-in approach to be viable it is necessary that the ‘pilot’ estimators to be plugged-in are accurate and not plagued by a difficult block/bandwidth choice themselves! Both of those issues can be successfully addressed by using pilot estimators of the trapezoidal lag-window type \hat{f}_{b,λ_T} ; see Paparoditis and Politis (2001a) or Politis and White (2001). Note that choosing b in the estimator \hat{f}_{b,λ_T} (or equivalently, in $\tilde{\sigma}_b^2$) is relatively straightforward: one may pick $b = 2\hat{m}$, where \hat{m} is the smallest integer such that $\hat{\gamma}(k) \simeq 0$ for $k > \hat{m}$. Here $\hat{\gamma}(k) \simeq 0$ really means $\hat{\gamma}(k)$ not significantly different from 0, i.e., an implied hypothesis test; see Politis (2001b) for more details.

Finally, note that all aforementioned block resampling/subsampling methods were designed for stationary time series. What happens if the time series is not stationary? Although both subsampling and block resampling will not break down under a mild nonstationarity—see e.g. Politis et al. (1999, Ch. 4)—the following interesting cases may be explicitly addressed:

- (i) *Seasonal effects.* If the time series X_t can be decomposed as $X_t = Y_t + W_t$ where Y_t is stationary and W_t is random or deterministic of period d , then block resampling and subsampling will generally remain valid if b is chosen to be an integer multiple of d ; in addition, resampling/subsampling should not be done from the set $\{B_1, B_2, \dots, B_{n-b+1}\}$ but rather from a set of the type $\{B_1, B_{d+1}, B_{2d+1}, \dots\}$ i.e., full overlap of the blocks is no longer recommended—see Politis (2001a).
- (ii) *Local stationarity.* Suppose that the time series has a slowly-changing stochastic structure in the sense that the joint probability law of $(X_t, X_{t+1}, \dots, X_{t+m})$ changes smoothly (and slowly) with t for any m ; see e.g. Dahlhaus (1997). Then, a ‘local’ block bootstrap may be advisable; here, a bootstrap pseudo-series is constructed by a concatenation of k blocks of size b (such that $kb \simeq n$), where the j th block

of the resampled series is chosen randomly from a distribution (say, uniform) on all the size- b blocks of consecutive data whose time indices are ‘close’ to those in the original block. A rigorous description of the local block bootstrap is given in Paparoditis and Politis (2002c).

- (iii) *Unit root-integrated time series.* A useful model in econometrics is a generalization of the random walk notion where X_t is not stationary but D_t is, where $D_t = X_t - X_{t-1}$ is the series of first differences; see e.g. Hamilton (1994). In that case, a block-bootstrap of the differences D_t can be performed yielding D_1^*, D_2^*, \dots , and a bootstrap pseudo-series for X_t can be constructed by ‘integrating’ the D_t^* , i.e., by letting $X_t^* = \sum_{i=1}^t D_i^*$. This idea is effectively imposing a sample-path ‘continuity’ on the bootstrap realization X_t^* ; see Paparoditis and Politis (2001c). On the other hand, if a unit root is just suspected to exist, a bootstrap test of the unit root hypothesis (vs. the alternative of stationarity) may be constructed using similar ideas; see Paparoditis and Politis (2000b).

2.4 Why have bootstrap methods been so popular?

Finally, it is interesting to recall the two main reasons for the immense success and popularity of the i.i.d. bootstrap of Efron (1979):

- (a) The bootstrap was shown to give valid estimates of distribution and standard error in ‘difficult’ situations; a prime example is the median of i.i.d. data for which the (delete-1) jackknife was known to fail, and the asymptotic distribution is quite cumbersome.
- (b) In ‘easy’ cases where there exist easy-to-construct alternative distribution estimators, e.g., the regular sample mean with its normal large-sample distribution, the bootstrap was shown to outperform those alternatives, i.e., to possess ‘second-order accuracy’—see Singh (1981) and Hall (1992).

In the case of dependent stationary data, under some regularity conditions, the appropriately² standardized/studentized block bootstrap (in all its variations including the stationary and circular bootstrap) also possesses a ‘higher-order accuracy’ property when estimating the distribution of the sample mean; see Lahiri (1991, 1999), and Götze and Künsch (1996). Nevertheless, the gains in accuracy are not as spectacular as in the i.i.d. case; in

²Note that it is not advisable to use a simple block bootstrap estimate of standard error for the studentization as it does not converge sufficiently fast; however, one may use an estimator such as $\tilde{\sigma}_b$ —see e.g. Bertail and Politis (2001).

addition, estimating the standard error accurately is the dominant factor so that it is somewhat of a luxury to think of capturing higher order moments.

The bottom line is that practically all time series problems fall under the ‘difficult’ category and thus the aforementioned block resampling/subsampling methods become invaluable. To give a simple example, consider the problem of estimating the distribution of the lag- p sample autocorrelation $\hat{\rho}(k) = \hat{\gamma}(k)/\hat{\gamma}(0)$, for some fixed k , with the purpose of constructing a confidence interval or hypothesis test concerning $\rho(k)$. It is quite straightforward to show, under usual moment and mixing conditions, that the large-sample distribution of $\sqrt{n}(\hat{\rho}(k) - \rho(k))$ is $N(0, V_k^2)$. Unfortunately, the variance V_k^2 is in general intractable involving infinite sums of fourth-order cumulants, etc.; see e.g. Romano and Thombs (1996). Nevertheless, block resampling and subsampling effortlessly yield consistent estimators of V_k^2 and the distribution of $\sqrt{n}(\hat{\rho}(k) - \rho(k))$.

One may ask how it had been possible in the pre-bootstrap days to practically address such a basic time series inference question regarding the autocorrelations; the answer is that people had to resort to some more restrictive model assumptions. For example, a celebrated formula for V_k^2 exists involving infinite sums of the (unknown) $\rho(\cdot)$ coefficients but no higher cumulants; this formula is due to Bartlett but is valid only under the assumption that the time series $\{X_t\}$ is *linear*, i.e., it satisfies an equation of the type

$$X_t = \mu + \sum_{i=-\infty}^{\infty} \psi_i Z_{t-i}, \quad (4)$$

where $Z_t \sim$ i.i.d. $(0, \tau^2)$, and the ψ_j coefficients are absolutely summable—see Brockwell and Davis (1991). Nonetheless, the class of *nonlinear* time series is vast; for example, the simple time series defined by $Y_t = Z_t Z_{t-1}$ for all t is nonlinear and 1-dependent although uncorrelated. Other examples of nonlinear time series models include the ARCH/GARCH and bilinear models that are of particular prominence in econometric analysis; see e.g. Engle (1995), Bollerslev et al. (1992), Granger and Andersen (1978), Subba Rao and Gabr (1984), or Tong (1990).

To give a concrete example, the $\pm 1.96/\sqrt{n}$ bands that most statistical programs like S+ automatically overlay on the correlogram (the plot of $\hat{\rho}(k)$ vs. k) are based on Bartlett’s formula. These bands are usually interpreted as giving an implied hypothesis test of the first few $\rho(k)$ being zero or not, $\rho(1)$ in particular; however, this interpretation is true *only* if one is willing to assume (4). If the time series $\{X_t\}$ is not linear, then Bartlett’s formula and the $\pm 1.96/\sqrt{n}$ bands are totally uninformative and may be misleading;

resampling and subsampling give the only practical solution in this case—see Romano and Thombs (1996) for more details.

3 Model-based resampling for time series

3.1 Linear autoregressions

Historically speaking, block resampling and/or subsampling were not the first approach to bootstrapping non-i.i.d. data. Almost immediately following Efron’s (1979) paper, the residual-based bootstrap for linear regression and autoregression was introduced and studied; cf. Freedman (1981, 1984), Efron and Tibshirani (1986, 1993). In the context of the semi-parametric $AR(p)$ model of eq. (1), the residual-based bootstrap amounts to an i.i.d. bootstrap of (a centered version of) the estimated residuals \hat{Z}_t defined by

$$\hat{Z}_t = (X_t - \bar{X}_n) - \hat{\phi}_1(X_{t-1} - \bar{X}_n) - \dots - \hat{\phi}_p(X_{t-p} - \bar{X}_n)$$

where $\hat{\phi}_j$ are some consistent estimators of ϕ_j , e.g., the Yule-Walker estimators. Having constructed bootstrap pseudo-residuals $\hat{Z}_1^*, \hat{Z}_2^*, \dots$, a bootstrap series X_1^*, \dots, X_n^* can be generated via the recursion (1) with $\hat{\phi}_j$ in place of ϕ_j ; finally, our statistic of interest can be recomputed from the bootstrap series X_1^*, \dots, X_n^* .

If the assumed $AR(p)$ model holds true, then the above residual-based bootstrap works very well for the sample mean and other more complicated statistics, and even enjoys a higher-order accuracy property similar to Efron’s i.i.d. bootstrap; see Bose (1988). Nevertheless, one never really knows whether the underlying stochastic structure conforms exactly to an $AR(p)$ model, and furthermore one would not know the value of p . A more realistic approach followed by a vast majority of practitioners is to treat eq. (1) only as an approximation to the underlying stochastic structure, and to estimate the order p from the data by criteria such as the AIC; see e.g. Shibata (1976). Doing so, one is implicitly assuming that $\{X_t\}$ is a linear time series possessing an $AR(\infty)$ representation of the form:

$$(X_t - \mu) = \sum_{i=1}^{\infty} \phi_i (X_{t-i} - \mu) + Z_t, \tag{5}$$

where $Z_t \sim \text{i.i.d. } (0, \tau^2)$, and the ϕ_j coefficients are absolutely summable.

Of course, in fitting model (5) to data, a finite order p must be used which, however, is allowed to increase as the sample size n increases. Thus,

the residual-based bootstrap applies *verbatim* to the AR(∞) model of eq. (5) with the understanding that p increases as a function of n ; this is the so-called AR(∞) “sieve” bootstrap employed by Swanepoel and van Wyk (1986), and rigorously investigated by Kreiss (1988, 1992), Paparoditis (1992), and Bühlmann (1997) among others. Conditions for validity of the AR(∞) bootstrap typically require that

$$p \rightarrow \infty \text{ as } n \rightarrow \infty \text{ but with } p^k/n \rightarrow 0 \quad (6)$$

where k is a small integer, usually 3 or 4. Consistency still holds when the order p is chosen via the AIC, i.e., it is data-dependent.

Under assumption (5), Choi and Hall (2000) proved a higher-order accuracy property of the AR(∞) bootstrap in the sample mean case, while Bühlmann (1997, 2002) showed that the AR(∞) bootstrap estimator $\hat{\sigma}_{AR(p)}^2$ of $\sigma_n^2 = \text{Var}(\sqrt{n}\bar{X}_n)$ has a performance comparable to the aforementioned optimal estimator $\tilde{\sigma}_b^2$. Both $\hat{\sigma}_{AR(p)}^2$ and $\tilde{\sigma}_b^2$ exhibit an *adaptivity* to the strength of dependence present in the series $\{X_t\}$, i.e., they become more accurate when the dependence is weak, and they both achieve a close to \sqrt{n} rate of convergence for series with exponentially decaying covariance $\gamma(s)$, e.g., in the case of stationary ARMA models. Not surprisingly, the best performance for $\hat{\sigma}_{AR(p)}^2$ is achieved when $\{X_t\}$ satisfies a finite-order AR model, i.e., a linear Markovian structure, whereas the best performance for $\tilde{\sigma}_b^2$ is achieved when $\{X_t\}$ has an m -dependent structure, e.g., a finite-order MA model. Nevertheless, $\tilde{\sigma}_b^2$ is more generally valid. Even among the class of linear time series (4) there are many series that do not satisfy (5), for example, the simple MA(1) model $X_t = Z_t + Z_{t-1}$, where $Z_t \sim \text{i.i.d.}(0, \tau^2)$.

3.2 Nonlinear autoregressions and Markov processes

Although the class of linear time series is quite rich including, e.g., all Gaussian time series, as mentioned before the class of nonlinear time series is vast. It is intuitive that the AR(∞) bootstrap will generally not give good results when applied to data X_1, \dots, X_n from a nonlinear time series³; see e.g. Bühlmann (2002) for a discussion and some illustrative examples.

The linearity of model (1) is manifested in the fact that the conditional expectation $m(X_{t-1}, \dots, X_{t-p}) := E(X_t | X_{t-1}, \dots, X_{t-p})$ is a linear function of the variables $(X_{t-1}, \dots, X_{t-p})$ just as in the Gaussian case. If we do not

³Unless of course it so happens that the statistic to be bootstrapped has a large-sample distribution that is totally determined by the first two moments of $\{X_t\}$, i.e., by μ and $\gamma(\cdot)$; this is actually what happens in the sample mean case.

know and/or assume that $m(\cdot)$ is linear, it may be of interest to estimate its functional form for the purposes of prediction of X_{n+1} given the recent observed past X_n, \dots, X_{n-p+1} . Under some smoothness assumptions on $m(\cdot)$, this estimation can be carried out by different nonparametric methods, e.g., kernel smoothing; see e.g. Masry and Tjøstheim (1995). The bootstrap is then called to provide a measure of accuracy of the resulting estimator $\hat{m}(\cdot)$, typically by way of constructing confidence bands. With this objective, many different resampling methods originally designed for nonparametric regression successfully carry over to this autoregressive framework. We mention some prominent examples:

- A. *Residual-based bootstrap*: Assuming the structure of a nonlinear (and heteroscedastic) autoregression:

$$X_t = m(X_{t-1}, \dots, X_{t-p}) + v(X_{t-1}, \dots, X_{t-p})Z_t, \quad (7)$$

where $Z_t \sim \text{i.i.d. } (0, 1)$, and $m(\cdot)$, $v(\cdot)$ are two (unknown) smooth functions, estimates of the residuals can be computed by $\hat{Z}_t = (X_t - \tilde{m}(X_{t-1}, \dots, X_{t-p}))/\tilde{v}(X_{t-1}, \dots, X_{t-p})$ where $\tilde{m}(\cdot)$, $\tilde{v}(\cdot)$ are preliminary (over-smoothed as in Härdle and Bowman (1988)) nonparametric estimators of $m(\cdot)$, $v(\cdot)$; then, an i.i.d. bootstrap can be performed on the (centered) residuals \hat{Z}_t , and recursion (7)—with $\hat{m}(\cdot)$, $\hat{v}(\cdot)$ in place of $m(\cdot)$, $v(\cdot)$ —can be used to generate a bootstrap series X_1^*, \dots, X_n^* from which our target statistic, e.g., $\hat{m}^*(\cdot)$, can be recomputed. The consistency of this procedure under some conditions has been recently shown in a highly technical paper by Franke et al. (2002).

- B. *Wild and Local bootstrap*: Let $\mathbf{X}_{t-1} = (X_{t-1}, \dots, X_{t-p})$ and consider the scatterplot of X_t vs. \mathbf{X}_{t-1} ; smoothing this scatterplot will give our estimator $\hat{m}(\cdot)$ as well as estimators for higher-order conditional moments if so desired. Interestingly, the wild bootstrap of Wu (1986) and the local bootstrap of Shi (1991) both have a direct application in this dependent context; their respective implementation can be performed exactly as in the independent (nonparametric) regression setting without even having to assume the special structure implied by the nonlinear autoregression (7) which is quite remarkable. Neumann and Kreiss (1998) show the validity of the wild bootstrap in this time series context, while Paparoditis and Politis (2000a) show the validity of the local bootstrap under very weak conditions.

Comparing the above two methods, i.e., the residual-based vs. the wild/local bootstrap we note the following: the wild/local bootstrap methods are valid

under weaker assumptions, e.g., (7) is not assumed; on the other hand, the residual-based bootstrap is able to generate new bootstrap stationary pseudo-series X_1^*, \dots, X_n^* from which a variety of statistics may be recomputed as opposed to just generating new scatterplots to be smoothed.

In order to have our cake and eat it too, the *Markovian local bootstrap* (MLB for short) was recently introduced by Paparoditis and Politis (2001b, 2002a). To describe it, assume for the sake of argument that the time series $\{X_t\}$ is Markov of order p , with $p = 1$ for simplicity. Given the data X_1, \dots, X_n , the MLB procedure constructs a bootstrap sample-path X_1^*, \dots, X_n^* as follows:

- (a) Assign X_1^* a value chosen randomly from the set $\{X_1, \dots, X_n\}$. Suppose this chosen value was equal to X_{t_1} (for some t_1).
- (b) Assign X_2^* a value chosen randomly from the set $\{X_{s+1} : |X_{t_1} - X_s| \leq h \text{ and } 1 \leq s < n - 1\}$, where h is a bandwidth-type parameter. Suppose this chosen value was equal to X_{t_2} .
- (c) Assign X_3^* a value chosen randomly from the set $\{X_{s+1} : |X_{t_2} - X_s| \leq h \text{ and } 1 \leq s < n - 1\}$. Repeat the above until the full bootstrap sample-path X_1^*, \dots, X_n^* is constructed.

The lag-1 (i.e., $p = 1$) MLB described above is capable of capturing the transition probability, as well as the two-dimensional marginal of the Markov process $\{X_t\}$; thus, it can capture the whole infinite-dimensional distribution of $\{X_t\}$ since that is totally determined by the two-dimensional marginal via the Markov property. To appreciate why, consider momentarily the case where $\{X_t\}$ has a finite state-space, say $\{1, 2, \dots, d\}$, and suppose that X_{t_1} from part (a) equals the state 1. Then we can even take $h = 0$, and the set $\{X_{s+1} : |X_{t_1} - X_s| \leq h\}$ is simply the set of all data points whose *predecessor* equaled 1. Choosing randomly from this set is like generating data from the empirical transition function, i.e., an estimate of $P(X_{t+1} = \cdot | X_t = 1)$. In the general continuous-state case, consistency of the MLB requires that we let $h \rightarrow 0$ as $n \rightarrow \infty$, but at a slow enough rate, e.g., $nh \rightarrow \infty$, sufficient to ensure that the cardinality of sets of the type $\{X_{s+1} : |X_t - X_s| \leq h\}$ increases with n .

The above MLB algorithm can be easily modified to capture the Markov(p) case for $p \geq 1$. It can then generally be proven—cf. Paparoditis and Politis (2001b, 2002a)—that the MLB bootstrap pseudo-series X_1^*, X_2^*, \dots is stationary and Markov of order p , and that the stochastic structure of $\{X_t^*\}$ accurately mimics the stochastic structure of the original Markov(p) series $\{X_t\}$.

Note that the assumption of a Markov(p) structure is more general than assuming eq. (7). It should be stressed, however, that the applicability of

the MLB is *not* limited to Markov processes: the lag- p MLB accurately mimics the $(p + 1)$ -joint marginal of a *general* stationary $\{X_t\}$ process, and therefore could be used for inference in connection with any statistic whose large-sample distribution only depends on this $(p + 1)$ -joint marginal; in that sense, the MLB is actually a *model-free* method. Prime examples of such statistics are the aforementioned kernel smoothed estimators of the conditional moments $m(\cdot)$ and $v(\cdot)$; see e.g. Paparoditis and Politis (2001b).

Finally note that a closely related method to the MLB is the Markov bootstrap of Rajarshi (1990) that also possesses many favorable properties; see e.g. Horowitz (2001) who also shows a higher-order accuracy property. The Markov bootstrap proceeds by nonparametrically estimating the transition density as a ratio of two kernel-smoothed density estimators (the joint over the marginal); then a bootstrap series X_1^*, \dots, X_n^* is generated by starting at an arbitrary data point, and then sampling from this explicitly estimated transition density.

The relation between the MLB and the Markov bootstrap of Rajarshi (1990) is analogous to the relation between Efron's (1979) i.i.d. bootstrap (that samples from the empirical) and the so-called smoothed bootstrap (that samples from a smoothed empirical); for example, the MLB only generates points X_t^* that actually belong to the set of data-points $\{X_1, \dots, X_n\}$ while the Markov bootstrap generates points X_t^* that can be anywhere on the real line. By analogy to the i.i.d. case it is intuitive here too that this extra smoothing may well be superfluous at least in the case of statistics such as the conditional/unconditional moments and functions thereof. Nevertheless, in different situations, for instance, if the statistic under consideration has a large-sample distribution that depends on the underlying marginal densities—as opposed to distributions—, this explicit smoothing step may be advisable. An example in the i.i.d. case is given by the sample median (and other non-extreme sample quantiles) where the smoothed bootstrap outperforms the standard bootstrap; see e.g. Hall et al. (1989).

References

- [1] Arcones, M.A. (2001). On the asymptotic accuracy of the bootstrap under arbitrary resampling size, submitted to *Ann. Instit. Statist. Math.*
- [2] Bartlett, M.S. (1946), On the theoretical specification of sampling properties of autocorrelated time series, *J.Roy.Statist.Soc.Suppl.*, 8, 27-41.

- [3] Bertail, P. and Politis, D.N. (2001). Extrapolation of subsampling distribution estimators: the i.i.d. and strong mixing cases, *Canadian J. Statistics*, 29, 667–680.
- [4] Bickel, P. and Freedman, D.A. (1981), Some asymptotic theory for the bootstrap, *Annals of Statistics*, 9, 1196-1217.
- [5] Bickel, P., Götze, F. and vanZwet, W.R. (1997), Resampling fewer than n observations: gains, losses, and remedies for losses, *Statistica Sinica*, 7, 1-32.
- [6] Bollerslev, T., Chou, R. and Kroner, K. (1992). ARCH modelling in finance: a review of theory and empirical evidence, *J. Econometrics*, 52, 5-60.
- [7] Booth, J.G. and Hall, P. (1993). An improvement of the jackknife distribution function estimator, *Annals of Statistics*, 21, 1476-1485.
- [8] Bose, A. (1988). Edgeworth correction by bootstrap in autoregressions, *Annals of Statistics*, 16, 1709-1722.
- [9] Brockwell, P. and Davis, R. (1991) *Time Series: Theory and Methods*, 2nd ed. Springer, New York.
- [10] Bühlmann, P. (1997), Sieve bootstrap for time series, *Bernoulli*, 3, 123-148.
- [11] Bühlmann, P. (2002). Bootstraps for time series. *Statistical Science*.
- [12] Carlstein, E. (1986). The use of subseries values for estimating the variance of a general statistic from a stationary time series. *Annals of Statistics* 14, 1171-1179.
- [13] Choi, E. and Hall, P. (2000). Bootstrap confidence regions computed from autoregressions of arbitrary order. *J. Roy. Statist. Soc., Ser. B*, 62, 461–477.
- [14] Dahlhaus, R. (1997). Fitting time series models to nonstationary processes, *Annals of Statistics*, 25, 1-37.
- [15] Efron, B. (1979). Bootstrap methods: another look at the jackknife, *Annals of Statistics*, 7, 1-26.

- [16] Efron, B. and Tibshirani, R.J. (1986). Bootstrap methods for standard errors, confidence intervals, and other measures of statistical accuracy (with discussion), *Statistical Science*, 1, 54-96.
- [17] Efron, B. and Tibshirani, R.J. (1993). *An Introduction to the Bootstrap*. New York: Chapman and Hall.
- [18] Engle, R. (1995). *ARCH Selected Readings*. Oxford: Oxford Univ. Press.
- [19] Franke, J., Kreiss, J.-P. and Mammen, E. (2002), Bootstrap of kernel smoothing in nonlinear time series, *Bernoulli*, Vol. 8, No. 1.
- [20] Freedman, D.A. (1981). Bootstrapping regression models, *Annals of Statistics*, 9, 1218-1228.
- [21] Freedman, D.A. (1984). On bootstrapping two-stage least-squares estimates in stationary linear models, *Annals of Statistics*, 12, 827-842.
- [22] Fuller, W.A. (1996). *Introduction to Statistical Time Series, 2nd ed.*, New York: John Wiley.
- [23] Giné, E. and Zinn, J. (1990). Necessary conditions for the bootstrap of the mean. *Annals of Statistics*, 17, 684–691.
- [24] Götze, F. and Künsch, H. (1996). Second order correctness of the blockwise bootstrap for stationary observations. *Annals of Statistics* 24, 1914-1933.
- [25] Granger, C. and Andersen, A. (1978). *An Introduction to Bilinear Time Series Models*, Göttingen: Vandenhoeck et Ruprecht.
- [26] Grenander, U. and Rosenblatt, M. (1957). *Statistical Analysis of Stationary Time Series*, New York: John Wiley.
- [27] Hall, P. (1985). Resampling a coverage pattern, *Stoch. Proc. Applic.*, 20, 231-246.
- [28] Hall, P. (1992). *The Bootstrap and Edgeworth Expansion*, New York: Springer.
- [29] Hall, P., DiCiccio, T. J. and Romano, J. P. (1989). On smoothing and the bootstrap. *Ann. Statist.* 17, 692–704.

- [30] Hall, P. Horowitz, J. L. and Jing, B.-Y. (1995), On Blocking Rules for the Bootstrap With Dependent Data, *Biometrika*, 82, 561-574.
- [31] Hamilton, J. D. (1994), *Time Series Analysis*, Princeton University Press, Princeton: New Jersey.
- [32] Härdle, W. and Bowman, A. (1988). Bootstrapping in nonparametric regression: local adaptive smoothing and confidence bands, *J. Amer. Statist. Assoc.*, 83, 100-110.
- [33] Horowitz, J.L. (2001). Bootstrap methods for Markov processes, Working paper, Dept. of Economics, Northwestern Univ., Evanston IL.
- [34] Kreiss, J. P. (1988). Asymptotic statistical inference for a class of stochastic processes. Habilitationsschrift, Faculty of Mathematics, University of Hamburg, Germany.
- [35] Kreiss, J. P. (1992). Bootstrap procedures for $AR(\infty)$. In *Bootstrapping and Related Techniques*, Joeckel, K. H., Rothe, G. and Sendler, W. (Eds.) Springer Verlag, Berlin, pp. 107-113
- [36] Künsch, H. R. (1989). The Jackknife and the Bootstrap for General Stationary Observations, *Annals of Statistics*, 17, 1217-1241.
- [37] Lahiri, S.N.(1991). Second order optimality of stationary bootstrap. *Statistics and Probability Letters* 11, 335-341.
- [38] Lahiri, S.N.(1999), Theoretical comparisons of block bootstrap methods, *Annals of Statistics*, 27, 386-404.
- [39] Liu, R.Y. and Singh, K. (1992). Moving blocks jackknife and bootstrap capture weak dependence. In *Exploring the limits of bootstrap*, ed. by LePage and Billard, pp. 225-248, New York: John Wiley.
- [40] Masry, E. and Tjøstheim, D. (1995). Nonparametric estimation and identification of nonlinear ARCH time series, *Econometric Theory*, 11, 258-289.
- [41] Neumann, M. and Kreiss, J.P. (1998). Regression-type inference in nonparametric autoregression. *Annals of Statistics*, 26, 1570-1613.
- [42] Paparoditis, E. (1992). Bootstrapping some statistics useful in identifying ARMA models. In *Bootstrapping and Related Techniques*, Joeckel, K. H., Rothe, G. and Sendler, W. (Eds.) Springer Verlag, Berlin, pp. 115-119.

- [43] Paparoditis, E. and Politis, D. N. (2000a), The Local Bootstrap for Kernel Estimators Under General Dependence Conditions, *Annals of Instit. Statist. Math.*, 52, 139-159.
- [44] Paparoditis, E. and Politis, D. N. (2000b), Residual-based Block Bootstrap for Unit Root Testing. Submitted to *Econometrica* in 2000.
- [45] Paparoditis, E. and Politis, D. N. (2001a). Tapered block bootstrap. *Biometrika* 88, 1105-1119.
- [46] Paparoditis, E. and Politis, D. N. (2001b). A Markovian local resampling scheme for nonparametric estimators in time series analysis. *Econometric Theory* 17, 540-566.
- [47] Paparoditis, E. and Politis, D. N. (2001c). The Continuous-Path Block-Bootstrap. In *Asymptotics in Statistics and Probability*, (Madan Puri, Ed.), pp. 305-320, Zeist (The Netherlands): VSP publications.
- [48] Paparoditis, E. and Politis, D. N. (2002a). The local bootstrap for Markov processes. *J. Statistical Planning and Inference*, vol. 108.
- [49] Paparoditis, E. and Politis, D. N. (2002b). The Tapered Block Bootstrap for General Statistics from Stationary Sequences, *Econometrics Journal*.
- [50] Paparoditis, E. and Politis, D. N. (2002c). Local block bootstrap, preprint.
- [51] Politis, D. N. (2001a). Resampling time series with seasonal components, in *Frontiers in Data Mining and Bioinformatics: Proceedings of the 33rd Symposium on the Interface of Computing Science and Statistics*, Orange County, California, June 13-17, 2001.
- [52] Politis, D. N. (2001b). Adaptive bandwidth choice. Submitted to *J. Nonparametric Statistics* in 2001.
- [53] Politis, D. N., and Romano, J.P. (1992a). A general resampling scheme for triangular arrays of α -mixing random variables with application to the problem of spectral density estimation. *Annals of Statistics* 20, 1985–2007.
- [54] Politis, D. N., and Romano, J.P. (1992b). A Circular Block-Resampling Procedure for Stationary Data. In *Exploring the Limits of Bootstrap*, (R. LePage and L. Billard, editors), pp. 263–270, New York: John Wiley.

- [55] Politis, D. N., and Romano, J.P. (1992c). A General Theory for Large Sample Confidence Regions Based on Subsamples under Minimal Assumptions, Tech. Report 399, Dept. of Statistics, Stanford University.
- [56] Politis, D. N., and Romano, J.P. (1993). Estimating the Distribution of a Studentized Statistic by Subsampling, *Bulletin of the International Statistical Institute, 49th Session*, Book 2, pp. 315-316.
- [57] Politis, D. N., and Romano, J.P. (1994a). The stationary bootstrap. *Journal of the American Statistical Association* 89, 1303–1313.
- [58] Politis, D. N., and Romano, J.P. (1994b). Large Sample Confidence Regions Based on Subsamples under Minimal Assumptions. *Annals of Statistics* 22, 2031-2050.
- [59] Politis, D. N., and Romano, J.P. (1995). Bias-Corrected Nonparametric Spectral Estimation, *J. Time Ser. Anal.*, 16, 67-104.
- [60] Politis, D. N., Romano, J.P. and Wolf, M. (1999). *Subsampling*. Springer Verlag, New York.
- [61] Politis, D. N., and White, H. (2001). Automatic block-length selection for the dependent bootstrap. Submitted to *Econometric Reviews*.
- [62] Quenouille, M. (1949). Approximate tests of correlation in time series. *J. Roy. Statist. Soc., Ser. B*, 11, 68-84.
- [63] Quenouille, M. (1956). Notes on bias in estimation. *Biometrika*, 43, 353-360.
- [64] Radulovic, D. (1996), The bootstrap of the mean for strong mixing sequences under minimal conditions, *Statist. Prob. Letters*, 28, 65-72.
- [65] Rajarshi, M.B. (1990). Bootstrap in Markov sequences based on estimates of transition density, *Annals of Instit. Statist. Math.*, 42, 253-268.
- [66] Romano, J.P. and Thombs, L. (1996). Inference for Autocorrelations under Weak Assumptions, *J. Amer. Statist. Assoc.*, 91, 590-600.
- [67] Sakov, A. and Bickel, P. (2000). An Edgeworth expansion for the m out of n bootstrapped median. *Statist. Prob. Lett.*, 49, 217-223.
- [68] Singh, K. (1981). On the asymptotic accuracy of Efron's bootstrap. *Annals of Statistics*, 9, 1187-1195.

- [69] Shao, J. and Wu, C.J.F. (1989). A general theory for jackknife variance estimation, *Annals of Statistics*, 17, 1176-1197.
- [70] Sherman, M. and Carlstein, E. (1996). Replicate histograms. *Journal of the American Statistical Association*, 91, 566-576.
- [71] Shi, S.G. (1991). Local bootstrap. *Annals Inst. Statist. Math.*, 43, 667-676.
- [72] Shibata, R. (1976). Selection of the order of an autoregressive model by Akaike's information criterion, *Biometrika*, 63, 117-126.
- [73] Subba Rao, T. and Gabr, M. (1984). *An Introduction to Bispectral Analysis and Bilinear Time Series Models*, Springer Lecture Notes in Statistics No. 24, New York.
- [74] Swanepoel, J.W.H. (1986). A note on proving that the (modified) bootstrap works. *Comm. Statist. Theory Methods*, 15, 3193-3203.
- [75] Swanepoel, J.W.H. and van Wyk, J.W.J. (1986). The bootstrap applied to spectral density estimation. *Biometrika*, 73, 135-142.
- [76] Tong, H. (1990). *Non-linear Time Series Analysis: a Dynamical Systems Approach*, Oxford Univ. Press, Oxford.
- [77] Tukey, J.W. (1958). Bias and confidence in not quite large samples (abstract). *Annals of Math. Statistics*, 29, 614.
- [78] Wu, C.J.F. (1986). Jackknife, bootstrap and other resampling methods in regression analysis (with discussion), *Annals of Statistics*, 14, 1261-1350.