

# Bi-graded Koszul modules, K3 carpets, and Green's conjecture

Steven Sam

University of California, San Diego

January 7, 2021

- Fix alg. closed field  $\mathbf{k}$  of characteristic 0.
- $C$  is a (smooth) genus  $g \geq 2$  curve with canonical bundle  $\omega_C$ .
- The canonical ring  $\Gamma_C = \bigoplus_{d \geq 0} H^0(C; \omega_C^{\otimes d})$  is finitely generated over  $A = \text{Sym } H^0(C; \omega_C) \cong \mathbf{k}[x_1, \dots, x_g]$ .
- We're concerned with vanishing of Betti numbers

$$\beta_{i,j}(C) = \dim_{\mathbf{k}} \text{Tor}_i^A(\Gamma_C, \mathbf{k})_j.$$

- **Green's conjecture** states that  $\beta_{i,i+2}(C) = 0$  for  $i < \text{Cliff}(C)$ , the Clifford index of  $C$ . This governs for how many steps the equations of  $C$  have only linear syzygies.
- For “most” curves,  $\text{Cliff}(C) = \text{gon}(C) - 2$  where  $\text{gon}(C)$  is the minimum degree of a non-constant map  $C \rightarrow \mathbf{P}^1$ .

- Voisin (2002, 2005): There is a nonempty Zariski dense set of curves in the moduli space of curves for which Green's conjecture holds. Geometric proof involving K3 surfaces.  
Note:  $\text{Cliff}(C) \leq (g - 1)/2$  for all  $C$   
Refinement: In fact, this set contains curves of each gonality.
- Aprodu–Farkas (2011): Green's conjecture holds for any curve that lies on a K3 surface.
- Many other variations...
- Aprodu–Farkas–Papadima–Raicu–Weyman (2019): Reproved Voisin's result using representation theory ideas (next slide).  
Method of proof is simpler and extends to positive characteristic  $p \geq (g + 2)/2$
- Schreyer (1986): Green's conjecture fails in low characteristic

- Betti numbers are semicontinuous, i.e., the Betti numbers of a flat degeneration can only go up. Hence, to prove (generic) vanishing, it suffices to prove it for a single example, and we can consider singular (smoothable) curves.
- A rational curve with  $g$  cusps has genus  $g$  and can be realized as a hyperplane section of the tangential surface  $T_g$  of the  $g$ -uple rational normal curve (= the union of its tangent lines).
- There is a short exact sequence of graded modules

$$0 \rightarrow \mathbf{k}[T_g] \rightarrow \widetilde{\mathbf{k}[T_g]} \rightarrow \omega_{\mathbf{k}[\mathbf{P}^1, \mathcal{O}(g)]}(-1) \rightarrow 0,$$

consisting of the homog. coordinate ring of  $T_g$ , its normalization, and the canonical module of the homog. coordinate ring of the  $g$ -uple RNC.

- The latter two can be understood, so it amounts to understanding a long exact sequence on Tor.

- The problem reduces to showing that the following map is surjective for  $i \leq (g-1)/2$ :

$$\begin{array}{ccc}
 \mathrm{Tor}_i^A(\widetilde{\mathbf{k}[T_g]}, \mathbf{k})_{i+1} & \longrightarrow & \mathrm{Tor}_i^A(\omega_{\mathbf{k}[P^1, \mathcal{O}(g)]}, \mathbf{k})_i \\
 \parallel & & \parallel \\
 \bigwedge^{i+1} (\mathrm{Sym}^{g-2} \mathbf{k}^2) \otimes D^{2i}(\mathbf{k}^2) & & \bigwedge^i (\mathrm{Sym}^{g-1} \mathbf{k}^2) \otimes \mathrm{Sym}^{g-2-i}(\mathbf{k}^2)
 \end{array}$$

- The group  $SL_2(\mathbf{k})$  acts on everything in sight, so the map is equivariant. However, it is difficult to guess a formula.
- There is more structure though: we fix  $i$  and sum over all  $g$ . It turns out that both are f.g. modules over  $\mathrm{Sym}(D^{i+1}\mathbf{k}^2)$  and the sum of maps is linear.
- The actual way forward is technical but insight comes from Hermite reciprocity isomorphism:

$$\mathrm{Sym}^n(D^m\mathbf{k}^2) \cong \bigwedge^m (\mathrm{Sym}^{m+n-1} \mathbf{k}^2).$$

- The key to using the module structure on the sum is that the cokernel can be recast as a **Koszul module**.
- Given a subspace  $K \subset \bigwedge^2 V$ , the Koszul module  $W(V, K)$  is the middle homology of the modified Koszul complex

$$\mathrm{Sym} V \otimes K \rightarrow \mathrm{Sym} V \otimes V(1) \rightarrow \mathrm{Sym} V(2)$$

- In previous setting,  $V = D^{i+1}\mathbf{k}^2$  and  $K = D^{2i}\mathbf{k}^2$ .
- AFPRW proved the following are equivalent:
  - $K^\perp \subset \bigwedge^2 V^*$  contains no nonzero rank 2 matrix
  - $W(V, K)$  is finite length
  - $W(V, K)_d = 0$  for all  $d \geq \dim V - 3$

This is enough to prove Green's conjecture for rational cuspidal curves, and hence for a nonempty dense subset of curves in the moduli space of curves.

- Double structures on  $\mathbf{P}^1$  (ribbons) give a different degeneration of genus  $g$  curves (intuition: if a non-hyperelliptic curve degenerates to a hyperelliptic one, this is the degeneration of the image of the canonical map)
- They are hyperplane sections of double structures on rational normal scrolls. More precisely, consider the projective space

$$\mathbf{P}^g = \mathbf{P}(\mathrm{Sym}^a \mathbf{k}^2 \oplus \mathrm{Sym}^{g-1-a} \mathbf{k}^2)$$

with an  $a$ -uple RNC and  $(g - 1 - a)$ -uple RNC. Let  $B$  be the homog. coordinate ring of the corresponding scroll.

- There is an extension

$$0 \rightarrow \omega_B \rightarrow B' \rightarrow B \rightarrow 0$$

where  $B'$  is the homog. coordinate ring of a **K3 carpet**. It is a double structure on the scroll.

- The ribbons are smoothable to curves of gonality  $a$ . Hence proving Green's conjecture for each ribbon proves it for most curves of each gonality (not just the maximal value ones).
- Can prove it holds in characteristic  $p \geq a$ . In generic case  $a = (g - 1)/2$ , this beats  $(g + 2)/2$  from cuspidal curves and resolves a conjecture of Eisenbud–Schreyer.
- Coordinate ring  $A$  of  $\mathbf{P}(\mathrm{Sym}^a \mathbf{k}^2 \oplus \mathrm{Sym}^{g-1-a} \mathbf{k}^2)$  is bigraded.
- The syzygies of  $\omega_B$  and  $B$  are understood, so we again need to consider a long exact sequence. The problem reduces to showing that the following map is surjective for  $i < a$ :

$$\begin{array}{ccc}
 \mathrm{Tor}_{i+1}^A(B, \mathbf{k})_{i+2} & \longrightarrow & \mathrm{Tor}_i^A(\omega_B, \mathbf{k})_{i+2} \\
 \parallel & & \parallel \\
 D^{i-1} \mathbf{k}^2 \otimes & & S^{g-3-i} \mathbf{k}^2 \otimes \\
 \bigwedge^{i+1} (S^{a-1} \mathbf{k}^2 \oplus S^{g-2-a} \mathbf{k}^2) & & \bigwedge^i (S^{a-1} \mathbf{k}^2 \oplus S^{g-2-a} \mathbf{k}^2)
 \end{array}$$



- We can decompose the last map into bigraded components, fix them, and sum over all  $a, g$ . Again, both terms are f.g. modules over a symmetric algebra and the cokernel is a bigraded Koszul module.
- Given vector spaces  $V_1, V_2$  and  $K \subset V_1 \otimes V_2 \subset \Lambda^2(V_1 \oplus V_2)$ ,  $W(V, K)$  is the middle homology of

$$\text{Sym}(V_1 \oplus V_2) \otimes K \rightarrow \begin{array}{c} \text{Sym}(V_1 \oplus V_2) \otimes V_1(0, 1) \\ \text{Sym}(V_1 \oplus V_2) \otimes V_2(1, 0) \end{array} \rightarrow \text{Sym}(V_1 \oplus V_2)$$

- In previous setting,  $V_1 = D^u \mathbf{k}^2$ ,  $V_2 = D^v \mathbf{k}^2$ ,  
 $K = D^{u+v-2} \mathbf{k}^2 + D^{u+v} \mathbf{k}^2$ .
- Raicu–Sam:
  - $K^\perp \subset V_1^* \otimes V_2^*$  contains no nonzero rank  $\leq 2$  matrix
  - $W(V, K)_{d,e} = 0$  for  $d, e \gg 0$
  - $W(V, K)_{d,e} = 0$  for  $d \geq \dim V_2 - 2$  and  $e \geq \dim V_1 - 2$ .

As before, this proves Green's conjecture for ribbons.