

Justify your answers! Put all the essential steps of your solution on this sheet!

1. Compute the equation of the tangent plane for the surface  $x^2y + yz = 6$  at the point  $(1,2,2)$ .
2. Is there a plane which contains the four points  $(1,1,0)$ ,  $(1,0,1)$ ,  $(0,0,2)$  and  $(2,2,0)$ ?
3. (a) In descending a hill whose equation is given by  $z = f(x, y) = 99 - x^2 + 2x - 1.5y^2$ , in which direction should a person standing at the point  $(3,1,94.5)$  initially proceed so as to descend as quickly as possible?  
(b) Is there any direction  $\mathbf{u}$  for which the directional derivative  $D_{\mathbf{u}}f(3, 1)$  is larger than 6? Why or why not?
4. Compute the double integral  $\int \int_D x dA$ , where  $D$  is the triangle with corners  $(0, 0)$ ,  $(1, 1)$  and  $(1, 0)$ .
5. Compute the integral  $\int \int_D e^{x^2+y^2} dA$ , where  $D$  consists of all points  $(x, y)$  with  $x^2 + y^2 \leq 9$  and  $x \geq 0$ .
6. Compute the surface area of the hyperbolic paraboloid  $z = x^2 - y^2$  which lies between the cylinders  $x^2 + y^2 = 1$  and  $x^2 + y^2 = 4$ .
7. Find the maximum and minimum of the function  $f(x, y, z) = 2x - z$  on the ellipsoid given by  $x^2 + 10y^2 + z^2 = 5$ , using Lagrange multipliers.
8. Compute max/min and saddle points, if any, for the function  $f(x, y) = 2x^3 + xy^2 + 5x^2 + y^2$ .
9. (a) Describe the part of the ball of radius 2 and center at the origin which lies below the cone  $z = -\sqrt{x^2 + y^2}$  in spherical coordinates.  
(b) Let  $f(x, y, z) = x^3 + \sin yz$  and let  $\rho, \theta, \phi$  be spherical coordinates. Find an expression of  $\partial f / \partial \phi$ . (It may contain ordinary as well as spherical coordinates).
10. Set up the triple integral which computes the volume of the solid between the plane  $z - 2x - y = 4$  and the paraboloid  $z = x^2 + y^2 + y$ . You need not compute the integral.

Justify your answers! Put all the essential steps of your solution on this sheet!

1. Compute the equation of the tangent plane for the surface  $x^2y + yz = 6$  at the point  $(1,2,2)$ .
2. Compute the integral  $\int \int_D \sin(x^2 + y^2) dA$ , where  $D$  consists of all points  $(x, y)$  with  $x^2 + y^2 \leq 9$  and  $x \geq 0$ .
3. Compute the double integral  $\int \int_D x dA$ , where  $D$  is the triangle with corners  $(0, 0)$ ,  $(1, 1)$  and

$(0, 1)$ .

4. (a) In descending a hill whose equation is given by  $z = f(x, y) = 99 - x^2 + 2x - 1.5y^2$ , in which direction should a person standing at the point  $(3, 1, 94.5)$  initially proceed so as to descend as quickly as possible?  
(b) Is there any direction  $\mathbf{u}$  for which the directional derivative  $D_{\mathbf{u}}f(3, 1)$  is larger than 6? Why or why not?
5. Compute the surface area of the hyperbolic paraboloid  $z = x^2 - y^2$  which lies between the cylinders  $x^2 + y^2 = 1$  and  $x^2 + y^2 = 4$ .
6. Is there a plane which contains the four points  $(1, 0, 1)$ ,  $(0, 2, 2)$ ,  $(2, 0, 0)$  and  $(0, 1, 1)$ ?
7. Find the maximum and minimum of the function  $f(x, y, z) = x - 2z$  on the ellipsoid given by  $x^2 + 10y^2 + z^2 = 5$ , using Lagrange multipliers.
8. Set up the triple integral which computes the volume of the solid between the plane  $z - x - 2y = 4$  and the paraboloid  $z = x^2 + y^2 + x$ . You need not compute the integral.
9. Compute max/min and saddle points, if any, for the function  $f(x, y) = 2x^3 + xy^2 + 5x^2 + y^2$ .
10. (a) Describe the part of the ball of radius 3 and center at the origin which lies below the cone  $z = -\sqrt{x^2 + y^2}$  in spherical coordinates.  
(b) Let  $f(x, y, z) = y^3 + \cos xz$  and let  $\rho, \theta, \phi$  be spherical coordinates. Find an expression of  $\partial f / \partial \phi$ . (It may contain ordinary as well as spherical coordinates).